Preprint typeset in JHEP style - PAPER VERSION HRI/ST/1409

arXiv:1405.3653v2 [hep-th] 22 Jul 2014

Exact results on ABJ theory and the refined topological string

Masazumi Honda[a] **and Kazumi Okuyama**[b]

[a] *Harish-Chandra Research Institute,*
Chhatnag Road, Jhusi, Allahabad 211019, India
[b] *Department of Physics, Shinshu University,*
3-1-1 Asahi, Matsumoto 390-8621, Japan

masazumihonda@hri.res.in, kazumi@azusa.shinshu-u.ac.jp

Abstract: We study the partition function of the ABJ theory, which is the $\mathcal{N} = 6$ superconformal Chern-Simons matter theory with gauge group $U(N) \times U(N + M)$ and Chern-Simons levels $(k, -k)$. We exactly compute the ABJ partition function on a three sphere for various k, M and N via the Fermi gas approach. By using these exact data, we show that the ABJ partition function is completely determined by the refined topological string on local $\mathbb{P}^1 \times \mathbb{P}^1$, including membrane instanton effects in the M-theory dual.

Keywords: Matrix Models, M-Theory, Topological Strings, AdS-CFT Correspondence.

Contents

1. Introduction **2**

2. ABJ theory as a Fermi Gas **4**
2.1 Rewriting canonical partition function 4
2.1.1 For $M \leq |k|/2$ 5
2.1.2 For $M > |k|/2$ 6
2.1.3 Remarks on Seiberg-like duality 7
2.2 Grand canonical formalism 7

3. Test of AdS/CFT correspondence **9**
3.1 Expectation from the gravity side 9
3.1.1 Classical supergravity 9
3.1.2 One-loop quantum supergravity 9
3.1.3 Non-perturbative effects 10
3.2 Comparison with the gravity side 10

4. ABJ partition function from the refined topological string **12**
4.1 Perturbative part 13
4.2 Non-perturbative effects and the refined topological string 14
4.2.1 The refined topological string 15
4.2.2 ABJ partition function 16
4.2.3 Cancellation of divergence between instantons 19
4.3 Test of our proposal 21
4.4 Comments on Matsumoto-Moriyama proposal 23

5. Discussions **24**

A. Algorithm for exact computation of ABJ grand partition function **24**
A.1 Algorithm for even Chern-Simons level 26
A.2 Algorithm for odd Chern-Simons level 27

B. π^{-N} term of $\widehat{Z}^{(N,N+M)}(k)$ **27**
B.1 $(k, M) = (2, 1)$ 28
B.2 $(k, M) = (4, 1)$ 28
B.3 Summary of the Higher Transcendental Part of Grand Partition Functions 29

C. Picard-Fuchs equation and effective Kähler parameters **29**

D. Exact results on ABJ partition function **30**

1. Introduction

Progress in M-theory would hinge on deeper understanding of its non-perturbative effects. A part of these effects appear as membrane instantons in M-theory [1]. During the last couple of years, there has been remarkable progress in understanding the membrane instanton effects. In [2], the authors have completely determined the non-perturbative structure of the free energy of M-theory on $AdS_4 \times S^7/\mathbb{Z}_k$ based on many different previous results. First of all, the M-theory on this background is expected to be dual to low-energy effective theory of multiple M2-branes [3]. Meanwhile it turns out that the M2-brane theory is described by an $\mathcal{N} = 6$ supersymmetric Chern-Simons matter theory (CSM) with gauge group $U(N)_k \times U(N)_{-k}$ commonly referred to as ABJM theory [4]. While M-theory region ($k \ll N^{1/5}$) cannot be accessed by the ordinary perturbative technique in the ABJM theory, the supersymmetry localization [5] reduces the partition function of the ABJM theory on S^3 to a matrix integral [6] called ABJM matrix model. This matrix model has been extensively studied by many approaches [7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 2, 20]. After the studies of the 't Hooft expansion [8, 10, 11, 7] and the leading order in the M-theory limit [9], there appeared a seminal work [13], which rewrites the ABJM matrix model as an ideal Fermi gas system (see also [12, 21]). This Fermi gas approach enables us to reveal structures of the partition function [15, 16, 17, 18, 19, 2, 20] and BPS Wilson loops [22, 23, 24] in the ABJM theory, including non-perturbative effects expected from the gravity side [25, 10, 1]. As a result, it turns out that the ABJM free energy is completely determined by the refined topological string on the "diagonal" local $\mathbb{P}^1 \times \mathbb{P}^1$ whose Kähler parameters for two $\mathbb{P}^1$'s are equal [2].

In this paper we extend the above analysis to the so-called ABJ theory [26, 27], which is the $\mathcal{N} = 6$ CSM with more general gauge group $U(N)_k \times U(N+M)_{-k}$. The ABJ theory is the low-energy effective theory of N M2-branes on $\mathbb{C}^4/\mathbb{Z}_k$ with M fractional M2-branes at the singularity. From the AdS/CFT perspective, one expects that this theory is dual to the M-theory on $AdS_4 \times S^7/\mathbb{Z}_k$ with a discrete torsion, and the type IIA superstring theory on $AdS_4 \times \mathbb{CP}^3$ with a nontrivial B-field holonomy. Furthermore the recent studies [28, 29] indicate that this theory has also a relation to the $\mathcal{N} = 6$ parity-violating Vasiliev theory on AdS_4 with a $U(N)$ gauge symmetry, when $M, k \gg 1$ with M/k and N kept fixed. Thus the ABJ theory is one of the most important supersymmetric gauge theories having M/string/higher-spin theory dual. The aim of this paper is to determine non-perturbative structure of free energy of the M-theory on $AdS_4 \times S^7/\mathbb{Z}_k$ with the discrete torsion via the supersymmetry localization, Fermi gas formalism, and refined topological string.

Let us briefly summarize our result[1]. Our starting point is a matrix integral representation for the ABJ partition on S^3 obtained by the localization method [6, 31, 32]:

$$Z^{(N,N+M)}(k) = \frac{i^{-\frac{1}{2}(N^2-(N+M)^2)\mathrm{sign}(k)}}{(N+M)!N!} \int_{-\infty}^{\infty} \frac{d^{N+M}\mu}{(2\pi)^{N+M}} \frac{d^N\nu}{(2\pi)^N} e^{-\frac{ik}{4\pi}(\sum_{j=1}^{N+M}\mu_j^2 - \sum_{a=1}^{N}\nu_a^2)}$$

[1] Recently an apparently similar conclusion was obtained in [30] by a different approach. While the values of the canonical partition function and finite part of grand potential presented in [30] are totally consistent with our values, there is an important difference of physical interpretation between [30] and ours. We will clarify an important difference between conclusions of [30] and our work in section 4.4.

$$\times \left[\frac{\prod_{1\le j<l\le N+M} 2\sinh\frac{\mu_j-\mu_l}{2} \prod_{1\le a<b\le N} 2\sinh\frac{\nu_a-\nu_b}{2}}{\prod_{j=1}^{N+M}\prod_{b=1}^{N} 2\cosh\frac{\mu_j-\nu_b}{2}} \right]^2, \tag{1.1}$$

which we call the ABJ matrix model. After conjectured in [33], the article [34] has proven that this matrix model has another equivalent representation suitable for a Fermi gas approach as explained in the next section. The Fermi gas expression of the ABJ matrix model is obtained by going to the grand canonical ensemble, *i.e.* by introducing the chemical potential μ and summing over N, as usual in the statistical mechanics. Moreover one can show that the computation of the grand canonical partition function boils down to construct a series of functions, which can be solved recursively. Thus we can exactly calculate the canonical partition function for various k, M, up to a fairly high $N = N_{\max}$. Indeed we have obtained exact values of the partition function for $(k, M, N_{\max}) = (2, 1, 65)$, $(3, 1, 31)$, $(4, 1, 62)$, $(4, 2, 29)$, $(6, 1, 23)$, $(6, 2, 21)$ and $(6, 3, 22)$, which are partly listed in appendix D. Then we compare these exact data with an expectation from the refined topological string on local $\mathbb{P}^1 \times \mathbb{P}^1$ with general Kähler parameters, which is a natural generalization of the ABJM case [2]. We will argue that the grand potential $J_k^{(M)}(\mu)$ of the ABJ theory is completely determined by the refined topological string free energy,

$$J_k^{(M)}(\mu) = F_{\rm top}(T_1^{\rm eff}, T_2^{\rm eff}; g_s) + \frac{1}{2\pi i}\frac{\partial}{\partial g_s}\left[g_s F_{\rm NS}\left(\frac{T_1^{\rm eff}}{g_s}, \frac{T_2^{\rm eff}}{g_s}; \frac{1}{g_s} \right) \right], \tag{1.2}$$

where the parameter g_s is related to the Chern-Simons level k as

$$g_s = \frac{2}{k}, \tag{1.3}$$

and $T_{1,2}^{\rm eff}$ are the effective Kähler parameters given by

$$T_1^{\rm eff}(\mu) = \frac{4\mu_{\rm eff}}{k} + 2\pi i\left(\frac{1}{2} - \frac{M}{k}\right), \quad T_2^{\rm eff}(\mu) = \frac{4\mu_{\rm eff}}{k} - 2\pi i\left(\frac{1}{2} - \frac{M}{k}\right), \tag{1.4}$$

with the effective chemical potential $\mu_{\rm eff}$ defined in (4.29). Here $F_{\rm top}$ and $F_{\rm NS}$ are the free energies of the un-refined topological string and the refined topological string in the Nekrasov-Shatashvili limit, respectively [35]. Although each individual terms in (1.2) have apparent divergences for physical integer k, a careful treatment shows that these divergences are actually canceled as in the ABJM case [19]. Then it remains a finite part, which will be presented in (4.39) for even k and (4.40) for odd k, respectively. Given the grand potential, we find that the ABJ partition function is written as the following sum of the Airy functions,

$$\begin{aligned} Z^{(N,N+M)}(k) = {} & C^{-\frac{1}{3}} e^{A+i\theta(N,M,k)} Z_{\rm CS}^{(M)}(k) \\ & \sum_{\ell,m=0}^{\infty} g_{\ell,m}\left(-\frac{\partial}{\partial N}\right) {\rm Ai}\left[C^{-\frac{1}{3}}\left(N - B + 2\ell + \frac{4m}{k} \right) \right], \end{aligned} \tag{1.5}$$

where $g_{\ell,m}$ is a polynomial explicitly determined by the grand potential $J_k^{(M)}(\mu)$, and C and B are the coefficients appearing in the perturbative part of the grand potential,

$$C = \frac{2}{\pi^2 k}, \quad B = \frac{1}{3k} - \frac{k}{12} + \frac{k}{2}\left(\frac{M}{k} - \frac{1}{2}\right)^2. \tag{1.6}$$

The overall constants $Z_{\rm CS}^{(M)}(k)$, $\theta(N,M,k)$ and A will be given by (2.2), (2.3) and (4.4), respectively. The indices (ℓ, m) label the number of D2-brane instantons and worldsheet instantons from the type IIA string viewpoint, while these are lifted up to membrane instantons wrapping two different three cycles in the M-theory. We will see that the large N asymptotic behavior of (1.5) reproduces the results of the classical supergravity (SUGRA) and the one-loop quantum SUGRA [36], and the correct weights of the instantons.

This paper is organized as follows. In section 2, we present the Fermi gas formalism of the ABJ partition function on S^3. In section 3, we compare the exact values of the partition function with the results on the gravity side. In section 4, we describe our conjecture (1.2) for the grand potential in terms of the refined topological string and perform a nontrivial test of (1.2) by using the exact data. Section 5 is devoted to discussion.

2. ABJ theory as a Fermi Gas

In this section we will show that the partition function of the ABJ theory on S^3 is described by an ideal Fermi gas system as in the ABJM case [13]. This picture enables us to compute the partition function exactly.

2.1 Rewriting canonical partition function

Let us consider the $U(N)_k \times U(N+M)_{-k}$ ABJ partition function on S^3. We begin with the ABJ matrix model (1.1) obtained by the localization method [6, 31, 32]. It is not so obvious (except $M = 0$ corresponding to the ABJM case) whether this expression of the ABJ partition function has an ideal Fermi gas description or not. However, it has been proven [34], after conjectured in [33], that the partition function has another equivalent representation, which is suitable for a Fermi gas approach. Although this representation takes seemingly different forms between $M \leq |k|/2$ and $M > |k|/2$, we will show below that the ABJ partition function can be rewritten for[2] all (k, M) as

$$Z^{(N,N+M)}(k) = e^{i\theta(N,M,k)} Z_{\rm CS}^{(M)}(k) \hat{Z}^{(N,N+M)}(k), \tag{2.1}$$

where $Z_{\rm CS}^{(M)}(k)$ is the $U(M)$ pure CS partition function on S^3 given by [39, 40, 41, 42]

$$Z_{\rm CS}^{(M)}(k) = |k|^{-\frac{M}{2}} \prod_{s=1}^{M-1} \left(2\sin\frac{\pi s}{|k|}\right)^{M-s}, \tag{2.2}$$

and $\theta(N,M,k)$ is the phase of the partition function,

$$e^{i\theta(N,M,k)} = i^{NM} e^{-\frac{i\pi}{6k}M(M^2-1)}. \tag{2.3}$$

The quantity $\hat{Z}^{(N,N+M)}(k)$ mainly used below, is the absolute value of the partition function divided by the pure CS contribution,

$$\hat{Z}^{(N,N+M)}(k) = \left| \frac{Z^{(N,N+M)}(k)}{Z_{\rm CS}^{(M)}(k)} \right|. \tag{2.4}$$

[2]For $M > |k|$, the ABJ partition function vanishes [33, 34] since $Z_{\rm CS}^{(M)}(k) = 0$ for this case. This is consistent with an expectation that spontaneously breaking of the supersymmetries occurs in this regime [26, 37, 38]. Hence we consider the case for $M \leq |k|$ below.

It turns out that $\hat{Z}^{(N,N+M)}(k)$ has a simple integral representation

$$\hat{Z}^{(N,N+M)}(k) = \frac{1}{N!}\int_{-\infty}^{\infty}\frac{d^N y}{(4\pi|k|)^N}\prod_{a<b}\tanh^2\frac{y_a-y_b}{2k}\prod_{a=1}^{N}V(y_a), \tag{2.5}$$

with

$$V(x) = \frac{1}{e^{\frac{x}{2}}+(-1)^M e^{-\frac{x}{2}}}\prod_{s=-\frac{M-1}{2}}^{\frac{M-1}{2}}\tanh\frac{x+2\pi i s}{2|k|}. \tag{2.6}$$

Here the product over s runs with the step $\Delta s = 1$. By using the Cauchy identity

$$\prod_{1\le a<b\le N}\tanh^2\frac{z_a-z_b}{2} = \sum_{\sigma\in S_N}(-1)^\sigma\prod_{a=1}^{N}\frac{1}{\cosh\frac{z_a-z_{\sigma(a)}}{2}},$$

we can rewrite the partition function as an ideal Fermi gas system,

$$\hat{Z}^{(N,N+M)}(k) = \frac{1}{N!}\sum_{\sigma\in S_N}(-1)^\sigma\int_{-\infty}^{\infty}\frac{d^N y}{(4\pi k)^N}\prod_{a=1}^{N}\rho(y_a,y_{\sigma(a)}), \tag{2.7}$$

where the density matrix $\rho(x,y)$ is given by

$$\rho(x,y) = \frac{\sqrt{V(x)V(y)}}{\cosh\frac{x-y}{2k}}. \tag{2.8}$$

In the rest of this section, we will prove that the ABJ partition function (1.1) can be written as (2.1) for all k and M.

2.1.1 For $M \le |k|/2$

For $M \le |k|/2$, the previous study has shown [34] that the ABJ partition function (1.1) is equivalent to

$$\begin{aligned} Z^{(N,N+M)}(k) = \frac{1}{N!}e^{i\theta(N,M,k)}Z_{\rm CS}^{(M)}(k)\int_{-\infty}^{\infty}\frac{d^N y}{(4\pi|k|)^N}\prod_{1\le a<b\le N}\tanh^2\frac{y_a-y_b}{2k} \\ \times\prod_{a=1}^{N}\left[\frac{1}{2\cosh\frac{y_a}{2}}\prod_{s=-\frac{M-1}{2}}^{\frac{M-1}{2}}\tanh\frac{y_a+2\pi i(s+M/2)}{2|k|}\right]. \end{aligned} \tag{2.9}$$

If we integrate this integrand along the contour $C_1+C_2+C_3+C_4$ depicted in fig. 1 (Left), then we can easily see that the integration vanishes by the Cauchy integration theorem[3]. Since the integrals along C_2 and C_4 becomes zero in the limit $\Lambda\to\infty$, we find

$$Z^{(N,N+M)}(k) = \int_{C_1}d^N y_a(\cdots) = \int_{C_3}d^N y_a(\cdots). \tag{2.10}$$

The right hand side is nothing but (2.1).

[3]Note that poles of $1/\cosh\frac{y_a}{2}$ at $y_a = -(2m+1)\pi i$ with $m = 1,2,\cdots,[(M-1)/2]$ are canceled out due to zeros of $\prod\tanh\frac{y_a+2\pi i(s+M/2)}{2k}$. Also, poles of $\tanh\frac{y_a+2\pi i(s+M/2)}{2|k|}$ are not located inside of $C_1+C_2+C_3+C_4$.



Figure 1: (Left) Explanation of the integral contours C_1, C_2, C_3 and C_4. The real positive parameter Λ is taken to infinity. (Right) Explanation of the integral contours C_5, C_6, C_7 and C_8.

2.1.2 For $M > |k|/2$

For $M > |k|/2$, the ABJ matrix model (1.1) is also written as [33, 34]

$$Z^{(N,N+M)}(k) = \frac{1}{N!} e^{i\theta(N,M,k)} Z_{\rm CS}^{(M)}(k) \int_{-\infty}^{\infty} \frac{d^N y}{(4\pi|k|)^N} \prod_{1\le a<b\le N} \tanh^2 \frac{y_a - y_b}{2k} \times \prod_{a=1}^{N} \left[\frac{1}{2\cosh\frac{y_a}{2}} \prod_{s=-\frac{|k|-M-1}{2}}^{\frac{|k|-M-1}{2}} \tanh \frac{y_a + 2\pi i(s + |k|/2 - M/2)}{2|k|} \right]. \quad (2.11)$$

By considering an integral contour $C_5 + C_6 + C_7 + C_8$ in fig. 1 (Right) with the same integrand, the partition function becomes

$$Z^{(N,N+M)}(k) = \int_{C_5} d^N y_a (\cdots) = \int_{C_7} d^N y_a (\cdots), \quad (2.12)$$

as in the case for $M \le |k|/2$. Thus we obtain

$$Z^{(N,N+M)}(k) = \frac{1}{N!} e^{i\theta(N,M,k)} Z_{\rm CS}^{(M)}(k) \int_{-\infty}^{\infty} \frac{d^N y}{(4\pi|k|)^N} \prod_{1\le a<b\le N} \tanh^2 \frac{y_a - y_b}{2k} \times \prod_{a=1}^{N} \left[\frac{1}{e^{\frac{y_a}{2}} + (-1)^{|k|-M} e^{-\frac{y_a}{2}}} \prod_{s=-\frac{|k|-M-1}{2}}^{\frac{|k|-M-1}{2}} \tanh \frac{y_a + 2\pi i s}{2|k|} \right]. \quad (2.13)$$

Noting that

$$\prod_{s=-\frac{|k|-1}{2}}^{\frac{|k|-1}{2}} \tanh \frac{y + 2\pi i s}{2|k|} = \begin{cases} 1 & \text{for } k : \text{even} \\ \tanh\frac{y}{2} & \text{for } k : \text{odd} \end{cases},$$

we find the following identity

$$\frac{1}{e^{\frac{y}{2}}+(-1)^{|k|-M}e^{-\frac{y}{2}}}\prod_{s=-\frac{|k|-M-1}{2}}^{\frac{|k|-M-1}{2}}\tanh\frac{y+2\pi is}{2|k|}=\frac{1}{e^{\frac{y}{2}}+(-1)^{M}e^{-\frac{y}{2}}}\prod_{s=-\frac{M-1}{2}}^{\frac{M-1}{2}}\tanh\frac{y+2\pi is}{2|k|}. \tag{2.14}$$

Plugging this into (2.13), we obtain the same expression (2.1).

2.1.3 Remarks on Seiberg-like duality

From the representation (2.1), one can see that the ABJ partition function transforms properly under the Seiberg-like duality [26] between the ABJ theories with different gauge groups

$$U(N)_k\times U(N+M)_{-k}\quad\text{and}\quad U(N+|k|-M)_k\times U(N)_{-k}. \tag{2.15}$$

This duality comes [43] from the Giveon-Kutasov duality [44], which has been proven for the S^3 partition function [45].

Let us check this duality in terms of (2.1). First, it is well-known that the pure CS partition function $Z_{\rm CS}^{(M)}(k)$ enjoys the level-rank duality (See e.g. [43] for a proof)

$$Z_{\rm CS}^{(M)}(k)=Z_{\rm CS}^{(|k|-M)}(-k). \tag{2.16}$$

Next, using (2.14) one can easily see that the integral representation of $\hat{Z}^{(N,N+M)}(k)$ in (2.5) is also duality invariant,

$$\hat{Z}^{(N,N+M)}(k)=\hat{Z}^{(N,N+|k|-M)}(-k). \tag{2.17}$$

Finally the phase factor $e^{i\theta(N,M,k)}$ given in (2.3) is not duality invariant but appropriately transforms as discussed in [43, 45].

In what follows, we will assume $k>0$ without loss of generality. Since $Z_{\rm CS}^{(M)}(k)$ and $\hat{Z}^{(N,N+M)}(k)$ depend only on the absolute value of k, (2.16) and (2.17) imply that they are invariant under the exchange $M\leftrightarrow k-M$:

$$Z_{\rm CS}^{(M)}(k)=Z_{\rm CS}^{(k-M)}(k),\qquad \hat{Z}^{(N,N+M)}(k)=\hat{Z}^{(N,N+k-M)}(k). \tag{2.18}$$

2.2 Grand canonical formalism

Let us switch to the grand canonical formalism. We define[4] the grand partition function as the generating function of $\hat{Z}^{(N,N+M)}(k)$ in (2.4) ,

$$\Xi_k^{(M)}(z)=\sum_{N=0}^{\infty}z^N\hat{Z}^{(N,N+M)}(k)=\sum_{N=0}^{\infty}z^N\left|\frac{Z^{(N,N+M)}(k)}{Z_{\rm CS}^{(M)}(k)}\right|, \tag{2.19}$$

[4]One could define the grand potential in terms of the whole partition function $Z^{(N,N+M)}(k)$ rather than $\hat{Z}^{(N,N+M)}(k)$. Then the N-dependent factor of the phase (2.3) can be absorbed by redefining the chemical potential as $\mu\to\mu+iM\pi/2$, while the N-independent factor of the phase and pure CS partition function $Z_{\rm CS}^{(M)}(k)$ can also be absorbed by redefinition of the perturbative coefficient A in (4.4). Particularly, in our definition of the grand partition function (2.19), we will see that corresponding topological string free energy is invariant under the Seiberg-like duality (2.15). If we did not drop the phase in defining the grand partition function, then corresponding topological string free energy do not become duality invariant.

where z is related to the chemical potential μ by

$$z = e^{\mu}. \tag{2.20}$$

From the representation (2.7) of $\hat{Z}^{(N,N+M)}(k)$ as the sum over permutations, one can show that the grand partition function is written as a Fredholm determinant,

$$\Xi_k^{(M)}(z) = \mathrm{Det}(1 + z\rho). \tag{2.21}$$

It is also convenient to introduce the grand potential as

$$\bar{J}_k^{(M)}(\mu) = \log \Xi_k^{(M)}(z). \tag{2.22}$$

Given the grand partition function $\Xi_k^{(M)}(z)$, we can easily come back to the canonical partition function $\hat{Z}^{(N,N+M)}(k)$ by

$$\hat{Z}^{(N,N+M)}(k) = \oint \frac{dz}{2\pi i} \frac{1}{z^{N+1}} \Xi_k^{(M)}(z). \tag{2.23}$$

As explained in [17, 46], the grand potential consists of a primary non-oscillatory part and an oscillatory part. The non-oscillatory part[5] $J_k^{(M)}(\mu)$ satisfies the following relation

$$e^{\bar{J}_k^{(M)}(\mu)} = \sum_{n=-\infty}^{\infty} e^{J_k^{(M)}(\mu + 2\pi i n)}. \tag{2.24}$$

Then we can deform the integral contour in (2.23) to the whole imaginary axis, by just replacing the total grand potential $\bar{J}_k^{(M)}(\mu)$ with its non-oscillatory part $J_k^{(M)}(\mu)$, namely,

$$\hat{Z}^{(N,N+M)}(k) = \int_{-i\infty}^{i\infty} \frac{d\mu}{2\pi i} \exp\Big[J_k^{(M)}(\mu) - \mu N\Big]. \tag{2.25}$$

Now let us describe our method for the exact computation of the partition functions. As discussed in [17] (see appendix A for details), thanks to the Tracy-Widom's lemma [47], the grand canonical partition function is determined by a series of functions $\phi_l^+(y)$,

$$\Xi_k^{(M)}(z) = \exp\left[-\sum_{n=1}^{\infty} \frac{z^{2n}}{n} \mathrm{Tr}\rho_+^{2n}\right] \cdot \sum_{l=0}^{\infty} \phi_l^+(0) z^l, \tag{2.26}$$

where ρ_+ and $\phi_l^+(y)$ are given by

$$\rho_+(x,y) = \frac{\rho(x,y) + \rho(x,-y)}{2} = \frac{E_+(x)E_+(y)}{\cosh\frac{x}{k} + \cosh\frac{y}{k}}, \quad E_+(x) = \cosh\frac{x}{2k}\sqrt{V(x)},$$

[5]The oscillatory part is defined by

$$\log\left[1 + \sum_{n=-\infty, n\neq 0}^{\infty} e^{J_k^{(M)}(\mu+2\pi i n) - J_k^{(M)}(\mu)}\right],$$

although we will not use this expression explicitly below.

$$\rho_+^n(x,y) = \frac{E_+(x)E_+(y)}{\cosh\frac{x}{k} + (-1)^{n-1}\cosh\frac{y}{k}} \sum_{l=0}^{n-1} (-1)^l \phi_l^+(x)\phi_{n-l-1}^+(y),$$

$$\phi_{l+1}^+(y) = \frac{1}{\cosh\frac{y}{2k}} \int \frac{dy'}{2\pi k} \frac{\cosh\frac{y'}{2k}}{2\cosh\frac{y-y'}{2k}} V(y')\phi_l^+(y'), \quad \phi_0^+(y) = 1,$$

$$\mathrm{Tr}\rho_+^{2n} = \int \frac{dy}{2\pi} \frac{E_+(y)^2}{\sinh\frac{y}{k}} \sum_{l=0}^{2n-1} (-1)^l \frac{d\phi_l^+(y)}{dy} \phi_{2n-l-1}^+(y). \tag{2.27}$$

In particular, $\phi_l^+(y)$ can be constructed recursively starting from $\phi_0^+(y) = 1$. Using the above formula, we have computed the exact partition functions of the ABJ theory for $(k, M, N_{\max}) = (2,1,65),\ (3,1,31),\ (4,1,62),\ (4,2,29),\ (6,1,23),\ (6,2,21)$ and $(6,3,22)$. The explicit values of the partition functions are listed in appendix D.

3. Test of AdS/CFT correspondence

In this section, using our exact data of the partition function, we test the AdS/CFT correspondence between the ABJ theory and M-theory on $AdS_4 \times S^7/\mathbb{Z}_k$ with a discrete torsion.

3.1 Expectation from the gravity side

Here we describe some expectations from the gravity side.

3.1.1 Classical supergravity

The $U(N)_k \times U(N+M)_{-k}$ ABJ theory [26, 27] describes the low-energy effective theory of N M2-branes with M fractional M2-branes. In the near-horizon limit, the geometry associated with this brane configuration becomes $AdS_4 \times S^7/\mathbb{Z}_k$ with M units of discrete torsion realized by a discrete holonomy of the 3-form field [26].

When $k \gg N^{1/5}$, the curvature becomes very small and we expect that the eleven-dimensional classical SUGRA on $AdS_4 \times S^7/\mathbb{Z}_k$ provides a good approximation of the gravity side. The free energy of the classical SUGRA with the boundary S^3 obeys the famous $N^{3/2}$-law [48], given by (see e.g. [49] for a derivation)

$$F_{\mathrm{SUGRA}} = -\frac{\pi\sqrt{2k}}{3} N^{3/2}. \tag{3.1}$$

In the next subsection we compare this with our exact result on the ABJ side in the large N regime with fixed k.

3.1.2 One-loop quantum supergravity

The one-loop correction of the SUGRA can also be analyzed by the technique successful in computing the logarithmic correction to the black hole entropy [50, 51, 52, 53, 54]. The authors in [36] have shown that the free energy of the eleven dimensional SUGRA

on $AdS_4 \times X_7$, where X_7 is a seven-dimensional manifold including $S^7/\mathbb{Z}_k$, contains the following universal logarithmic correction,

$$-\frac{1}{4}\log N. \tag{3.2}$$

For $\mathcal{N}=3$ necklace quiver CSM with coincident rank of gauge groups, it has been shown that this logarithmic behavior comes from the large N asymptotics of the Airy function [13] in the perturbative part of partition function (See also [55] for $\mathcal{N}=2$ case).

Strictly speaking, in the presence of the discrete torsion, there might be further massless degrees of freedom and the logarithmic correction could change[6]. However, as we will see in section 4, the exact ABJ partition functions show a nice agreement with the Airy function, and hence the ABJ partition function also exhibit this 1-loop behavior (3.2).

3.1.3 Non-perturbative effects

One expects two kinds of non-perturbative effects on the gravity side as in the ABJM case [25, 10]. If we identify a direction of M-theory circle with the orbifolding direction and consider a large k regime, then the eleventh dimension in the geometry $AdS_4 \times S^7/\mathbb{Z}_k$ shrinks and the bulk theory reduces to the type IIA string theory on $AdS_4 \times \mathbb{CP}^3$. Since $\mathbb{CP}^3$ has a nontrivial 2-cycle $\mathbb{CP}^1$ and a Lagrangian submanifold $\mathbb{RP}^3$, we expect that the dual type IIA string theory has the worldsheet instanton [25] and the D2-brane instanton [10] corrections characterized by the following weights, respectively,

$$\exp\left[-T_{\rm F1}{\rm Vol}(\mathbb{CP}^1)\right] = \exp\left(-2\pi\sqrt{\frac{2N}{k}}\right), \quad \exp\left[-T_{\rm D2}{\rm Vol}(\mathbb{RP}^3)\right] = \exp\left(-\pi\sqrt{2kN}\right). \tag{3.3}$$

As discussed in [8], the worldsheet instanton also receives the following contribution from the coupling to the background NSNS 2-form field

$$\int_{\mathbb{CP}^1} B_{\rm NS} = \frac{1}{2} - \frac{M}{k}. \tag{3.4}$$

We also expect that the D2-instanton has a coupling to the background RR 3-form field. From the viewpoint of M-theory, these instantons correspond to M2-branes wrapping three cycles known as membrane instantons [1]. The worldsheet instanton effects in (3.3) and (3.4) have been successfully reproduced from the ABJ matrix model in the 't Hooft limit [8]. In section. 4, we will show that the ABJ partition function contains all the expected instanton effects and determine the structure from the refined topological string.

3.2 Comparison with the gravity side

Let us compare our exact result of the partition function with the classical SUGRA result (3.1). In fig. 2 (Left), we plot the exact values of the free energy $\log \hat{Z}^{(N,N+M)}(k)$ listed in appendix D against $N^{3/2}$ for some cases. As expected from the gravity side, we observe that the free energy is approximately proportional to $N^{3/2}$ in the large N regime. Fig. 2

[6] We would like to thank Ashoke Sen for explaining this point.



Figure 2: (Left) The free energy $\log \hat{Z}^{(N,N+M)}(k)$ is plotted to $N^{3/2}$. (Right) The ratio between the exact free energy $\log \hat{Z}^{(N,N+M)}(k)$ and the classical SUGRA free energy $F_{\rm SUGRA}$ is plotted against $1/N$. The symbols and dashed lines denote the exact data and fitting functions in the large N region, respectively.

(Right) compares our result with the classical SUGRA result. In this plot, we fit the ratio between the exact free energy and classical SUGRA free energy by linear functions of $1/N$, whose intercepts should be almost 1 if the AdS/CFT correspondence is correct. We easily see that the data points are well fitted by the fitting functions in the large N region, and indeed the values of these intercepts are almost 1. Although we have plotted only for $(k, M) = (2, 1), (3, 1), (4, 2)$ and $(6, 3)$, we have checked that similar result holds also for $(k, M) = (4, 1), (6, 1)$ and $(6, 2)$.

We can reproduce the classical SUGRA result also from the thermodynamic limit of the Fermi gas system as in the ABJM case [13]. In the large N limit, the free energy is approximated by a saddle point of the μ-integration in (2.25) as

$$\log \hat{Z}^{(N,N+M)}(k) \simeq J_k^{(M)}(\mu_*) - \mu_* N, \qquad \text{with} \quad \left.\frac{\partial J_k^{(M)}(\mu)}{\partial \mu}\right|_{\mu=\mu_*} = N. \tag{3.5}$$

Hence we can estimate the large N free energy if we find the large μ asymptotic behavior of the grand potential, which is captured by a semi-classical expansion of the Fermi gas system. By introducing the quantum mechanical operators $\hat{q}$ and $\hat{p}$ satisfying $[\hat{q}, \hat{p}] = 2\pi i k$, and $\hat{q}|x\rangle = x|x\rangle$, we can rewrite the density matrix $\rho(x, y)$ as a quantum mechanical operator

$$\rho(x, y) = \langle x|e^{-\hat{H}(\hat{q},\hat{p})}|y\rangle, \quad e^{-\hat{H}(\hat{q},\hat{p})} = \sqrt{V(\hat{q})}\frac{1}{2\cosh\frac{\hat{p}}{2}}\sqrt{V(\hat{q})}, \tag{3.6}$$

where $\hat{H}$ is the Hamiltonian of the Fermi gas system. The asymptotic behavior of the classical Hamiltonian for $|q|, |p| \gg 1$ is given by

$$H_{\rm cl}(q, p) \sim \frac{|q| + |p|}{2}, \tag{3.7}$$

which is exactly the same as in the ABJM case [13]. Thus the same analysis as [13] gives the large μ grand potential and the saddle point μ_* as

$$J_k^{(M)}(\mu) \simeq \frac{2}{3\pi^2 k}\mu^3, \quad \mu_* = \pi\sqrt{\frac{kN}{2}}. \tag{3.8}$$

We can easily see that plugging this into (3.5) reproduces the result of the classical supergravity.

Using the expression of the saddle point value μ_* in (3.8), we can translate the instanton effects (3.3) on the gravity side into the language of the grand canonical formalism. In terms of μ_*, we can express the instanton factors as

$$\exp\left(-2\pi\sqrt{\frac{2N}{k}}\right) = \exp\left(-2\mu_*\right), \quad \exp\left(-\pi\sqrt{2kN}\right) = \exp\left(-\frac{4\mu_*}{k}\right). \tag{3.9}$$

If we assume that the free energy on the gravity side receives a series of multi-instanton corrections, we would expect that the ABJ grand potential have the following expansion,

$$J_k^{(M)}(\mu) = \sum_{l,m=0}^{\infty} f_{l,m}(\mu) \exp\left[-\left(2l + \frac{4m}{k}\right)\mu\right], \tag{3.10}$$

where $f_{l,m}(\mu)$ is a polynomial of μ, whose coefficients depend on M and k. Although there is no currently available technique to determine the coefficients $f_{l,m}(\mu)$ of instantons from the computation of the gravity side, the ABJ matrix model and its relation to the refined topological string give a very concrete prediction for these coefficients, as we will see in the next section.

4. ABJ partition function from the refined topological string

In this section we propose that the ABJ free energy including the non-perturbative effects is completely determined by the refined topological string on local $\mathbb{P}^1 \times \mathbb{P}^1$. As explained in [7], the partition function of the ABJ theory on S^3 can be analytically continued[7] to the partition function of the $L(2,1)$ lens space matrix model [39, 40] which comes from the pure Chern-Simons theory on $S^3/\mathbb{Z}_2$. The lens space matrix model corresponds to the topological string on local $\mathbb{P}^1 \times \mathbb{P}^1$ via a topological version of the large N duality [59]. Thus we expect that structure of the ABJ free energy is captured by the topological string on local $\mathbb{P}^1 \times \mathbb{P}^1$.

For the ABJM case, as the special case of the ABJ theory, this expectation has been confirmed quite successfully. In [2] it is discussed that the grand potential of the ABJM theory corresponds to the free energy of the topological string on the "diagonal" local $\mathbb{P}^1 \times \mathbb{P}^1$ in large radius frame, where the Kähler parameters of two $\mathbb{P}^1$'s are equal. More precisely, it has been shown, based on the exact and numerical results [15, 16, 17, 19], that the perturbative and worldsheet instanton part of the ABJM grand potential are given by the free energy of the un-refined topological string on the local $\mathbb{P}^1 \times \mathbb{P}^1$ [13, 17], while the D2-brane instanton part and its mixed contribution with the worldsheet instanton are captured by the Nekrasov-Shatashvili limit [35] of the refined topological string on the same local $\mathbb{P}^1 \times \mathbb{P}^1$.

In this paper we generalize this argument to the ABJ theory. From the topological string viewpoint, this amounts to consider non-diagonal local $\mathbb{P}^1 \times \mathbb{P}^1$, or equivalently local

[7] There is much strong evidence for this relation [7, 10, 14, 17, 33, 34, 56, 57, 58].

$\mathbb{P}^1 \times \mathbb{P}^1$ with general Kähler parameters. For this purpose, we decompose $J(\mu)$ into the perturbative part $J_{\rm pert}(\mu)$ and the non-perturbative part $J_{\rm np}(\mu)$ as

$$J_k^{(M)}(\mu) = J_{\rm pert}(\mu) + J_{\rm np}(\mu). \tag{4.1}$$

We propose that structure of the ABJ grand potential, or equivalently the ABJ partition function, is completely determined by the topological string including the non-perturbative effect and test this proposal by using our exact data of the partition functions.

4.1 Perturbative part

In order to study the non-perturbative structure of the ABJ grand potential in terms of the exact data, we shall first determine the perturbative part of the ABJ partition function to subtract this from the exact data. In this subsection, we determine the perturbative part of the ABJ grand potential by using the un-refined topological string.

As explained in [60], the perturbative part of the free energy of un-refined topological string on a Calabi-Yau manifold (CY) X is given by

$$F_{\rm pert} = \frac{1}{6g_s^2}\int_X J\wedge J\wedge J - \frac{1}{24}\left(1-\frac{1}{g_s^2}\right)\int_X J\wedge c_2, \tag{4.2}$$

where J is the Kähler form and c_2 is the second Chern class. From this general expression, we expect that the perturbative part of the grand potential of ABJ theory is given by

$$J_{\rm pert}(\mu) = \frac{T_1^3+T_2^3-3T_1^2T_2-3T_1T_2^2}{6g_s^2(4\pi i)^2} + \frac{1}{24}\left(1-\frac{1}{g_s^2}\right)(T_1+T_2) + A, \tag{4.3}$$

where $T_{1,2}$ are the Kähler parameters of local $\mathbb{P}^1\times\mathbb{P}^1$, whose relation to μ will be explained shortly. Here g_s denotes the string coupling $g_s = 2/k$ defined in (1.3) and A is given by

$$\begin{aligned} A = &-\log|Z_{\rm CS}^{(M)}(k)| - \frac{\zeta(3)k^2}{8\pi^2} + \frac{1}{6}\log\frac{4\pi}{k} + 2\zeta'(-1) \\ &- \frac{1}{3}\int_0^\infty dx\left(\frac{3}{x^3}-\frac{1}{x}-\frac{3}{x\sinh^2 x}\right)\frac{1}{e^{kx}-1}. \end{aligned} \tag{4.4}$$

Apart from the first term in (4.4), A is the so-called constant map contribution analyzed in [14] in detail. The first term $-\log|Z_{\rm CS}^{(M)}(k)|$ in (4.4) comes from our definition of the grand partition function (2.19). In order to reproduce the worldsheet instanton factor $e^{-4\mu/k}$ (3.9) together with the effect of the B-field (3.4), it is natural to make the following identification of the Kähler parameters

$$T_1(\mu) = \frac{4\mu}{k} + 2\pi i\left(\frac{1}{2}-\frac{M}{k}\right), \quad T_2(\mu) = \frac{4\mu}{k} - 2\pi i\left(\frac{1}{2}-\frac{M}{k}\right). \tag{4.5}$$

With this identification, the Seiberg-like duality (2.18) is naturally realized as the exchange of two $\mathbb{P}^1$'s of local $\mathbb{P}^1\times\mathbb{P}^1$

$$M \leftrightarrow k-M \quad \Leftrightarrow \quad T_1 \leftrightarrow T_2. \tag{4.6}$$

Then the perturbative grand potential in (4.3) becomes

$$J_{\rm pert}(\mu) = \frac{C}{3}\mu^3 + B\mu + A, \tag{4.7}$$

where C and B are defined in (1.6). This expression (4.7) of the perturbative part is consistent with the result for the ABJM case [13] and the same as the proposal in [30]. Then, as in the ABJM case, the perturbative canonical partition function $\hat{Z}^{(N,N+M)}_{\rm pert}(k)$ is given by the Airy function,

$$\hat{Z}^{(N,N+M)}_{\rm pert}(k) = \int_{-i\infty}^{i\infty} \frac{d\mu}{2\pi i} e^{J_{\rm pert}(\mu) - N\mu} = C^{-1/3} e^{A} {\rm Ai}[C^{-1/3}(N-B)]. \tag{4.8}$$

Below, we will show that this is indeed the correct perturbative partition function by comparing with our exact data. In the large N limit with fixed k, the perturbative part of the free energy becomes

$$\log \hat{Z}^{(N,N+M)}_{\rm pert}(k) = -\frac{2}{3}C^{-1/2}N^{3/2} + C^{-1/2}BN^{1/2} - \frac{1}{4}\log N + \mathcal{O}(1). \tag{4.9}$$

The first term reproduces the classical SUGRA result (3.1) and the third term agrees with the logarithmic behavior (3.2) of the one-loop quantum supergravity [36].

Let us compare $\hat{Z}^{(N,N+M)}_{\rm pert}(k)$ with our exact data of partition functions. For this purpose we decompose the canonical partition function as

$$\hat{Z}^{(N,N+M)}(k) = \hat{Z}^{(N,N+M)}_{\rm pert}(k)\left(1 + \hat{Z}^{(N,N+M)}_{\rm np}(k)\right), \tag{4.10}$$

where

$$\hat{Z}^{(N,N+M)}_{\rm np}(k) = \frac{\hat{Z}^{(N,N+M)}(k)}{\hat{Z}^{(N,N+M)}_{\rm pert}(k)} - 1. \tag{4.11}$$

If $\hat{Z}^{(N,N+M)}_{\rm pert}(k)$ is the correct perturbative part, then $\hat{Z}^{(N,N+M)}_{\rm np}(k)$ should correspond to the non-perturbative part of the partition function and it is expected to behave as

$$\hat{Z}^{(N,N+M)}_{\rm np}(k) = \mathcal{O}\Big(e^{-2\pi\sqrt{\frac{2N}{k}}}\Big), \tag{4.12}$$

in the large N regime. Fig. 3 shows the plots of $\hat{Z}^{(N,N+M)}_{\rm np}(k)$ against $2\pi\sqrt{2N/k}$ in semi-log scale for $(k,M) = (2,1), (6,1), (6,2)$ and $(6,3)$. We can easily see that the data points are on the straight lines in the large N regime, which imply the exponentially suppressed behavior of $\hat{Z}^{(N,N+M)}_{\rm np}(k)$ by $e^{-2\pi\sqrt{\frac{2N}{k}}}$. Thus we conclude that $J_{\rm pert}(\mu)$ given in (4.7) is indeed the correct perturbative part of the grand potential. Since this consistency check shows the presence of the "$-\frac{1}{4}\log N$" term (3.2) in the canonical partition function, we have also tested the AdS/CFT correspondence at 1-loop level of the dual quantum supergravity [36]. Although we have not explicitly shown, similar result holds also for $(k,M) = (3,1), (4,1)$ and $(4,2)$.

4.2 Non-perturbative effects and the refined topological string

In this section we briefly review [2] and write down an expected form of the ABJ grand potential from the refined topological string on local $\mathbb{P}^1 \times \mathbb{P}^1$ with non-diagonal Kähler parameters.



Figure 3: $\hat{Z}_{\rm np}^{(N,N+M)}(k)$ is plotted to $2\pi\sqrt{2N/k}$ in semi-log scale. (Left) A plot for $(k, M) = (2, 1)$. (Right) Plots for $(k, M) = (6, 1), (6, 2)$ and $(6, 3)$. The blue circle, purple square and yellow diamond symbols show the cases for $M = 1$, $M = 2$ and $M = 3$, respectively.

4.2.1 The refined topological string

Let us first recall the structure of the un-refined topological string on CY X with Kähler parameters T_I. The free energy of the un-refined topological string in the large radius frame is given by [61]

$$F_{\rm top}(T_I; g_s) = \sum_{g\geq 0}\sum_{w\geq 1}\sum_{\mathbf{d}} \frac{(-1)^{g-1}}{w} n_g^{\mathbf{d}} \left(q_s^{w/2} - q_s^{-w/2}\right)^{2g-2} \mathbf{Q}^{w\mathbf{d}}, \tag{4.13}$$

where

$$q_s = e^{\hat{g}_s}, \quad \mathbf{d} = (d_1, d_2, \cdots), \quad \mathbf{Q} = \prod_I Q_I, \quad Q_I = e^{-T_I}, \quad \mathbf{Q}^{\mathbf{d}} = \prod_I Q_I^{d_I}. \tag{4.14}$$

Here $\hat{g}_s$ is the topological string coupling defined by $\hat{g}_s = 2\pi i g_s$. The integer numbers $n_g^{\mathbf{d}}$ are the so-called Gopakumar-Vafa invariants of X.

The free energy of the refined topological string is obtained by a one parameter deformation of the standard topological string free energy originated in the Nekrasov instanton partition function [62]. The refined free energy is determined by the supersymmetric index $N_{j_L,j_R}^{\mathbf{d}}$ which counts the BPS states in the M-theory compactified on X down to 5 dimensions [63]. These BPS states come from M2-branes wrapping a two-cycle of X with degree $\mathbf{d}$. They behave as massive particles in the 5 dimensions, carrying spins (j_L, j_R) of the little group $SO(4) \simeq SU(2) \times SU(2)$. The refined free energy takes the form of

$$F_{\rm ref}(T_I; \epsilon_1, \epsilon_2) = \sum_{j_L,j_R\geq 0}\sum_{w\geq 1}\sum_{\mathbf{d}} \frac{(-1)^{s_L+s_R}}{w} N_{j_L,j_R}^{\mathbf{d}} \frac{\chi_{j_L}(q_L^w)\chi_{j_R}(q_R^w)}{(q_1^{w/2} - q_1^{-w/2})(q_2^{w/2} - q_2^{-w/2})} \mathbf{Q}^{w\mathbf{d}}, \tag{4.15}$$

where

$$s_{L,R} = 2j_{L,R} + 1, \quad q_L = e^{\frac{\epsilon_1-\epsilon_2}{2}}, \quad q_R = e^{\frac{\epsilon_1+\epsilon_2}{2}}, \quad q_{1,2} = e^{\epsilon_{1,2}}, \quad \chi_j(q) = \frac{q^{2j+1} - q^{-2j-1}}{q - q^{-1}}. \tag{4.16}$$

It is often useful to introduce another invariant $n^{\mathbf{d}}_{g_L.g_R}$ by

$$\sum_{j_L,j_R\geq 0}(-1)^{s_L+s_R}N^{\mathbf{d}}_{j_L,j_R}\chi_{j_L}(q_L)\chi_{j_R}(q_R)=\sum_{g_L,g_R\geq 0}n^{\mathbf{d}}_{g_L,g_R}(q_L^{\frac{1}{2}}-q_L^{-\frac{1}{2}})^{2g_L}(q_R^{\frac{1}{2}}-q_R^{-\frac{1}{2}})^{2g_R}. \quad (4.17)$$

The un-refined topological string corresponds to the following particular limit:

$$\epsilon_1=-\epsilon_2=\hat{g}_s, \quad (4.18)$$

with an identification

$$n^{\mathbf{d}}_g=n^{\mathbf{d}}_{g,0}. \quad (4.19)$$

As discussed in [2], the Nekrasov-Shatashvili (NS) limit

$$\epsilon_1=\hbar, \quad \epsilon_2\to 0. \quad (4.20)$$

of the refined topological string plays an important role in the D2-instanton corrections. Since the refined free energy has a simple pole ($\sim 1/\epsilon_2$) in this limit, we introduce the NS free energy as

$$\begin{aligned}F_{\rm NS}(T_I;\hbar) &= \lim_{\epsilon_2\to 0}\epsilon_2 F(T_I;\hbar,\epsilon_2)\\ &= \sum_{j_L,j_R\geq 1}\sum_{w\geq 1}\sum_{\mathbf{d}}\frac{(-1)^{s_L+s_R}}{w^2}N^{\mathbf{d}}_{j_L,j_R}\frac{\chi_{j_L}(q^{w/2})\chi_{j_R}(q^{w/2})}{q^{w/2}-q^{-w/2}}\mathbf{Q}^{w\mathbf{d}}, \end{aligned} \quad (4.21)$$

where

$$q=e^{\hbar}=e^{2\pi i/g_s}. \quad (4.22)$$

4.2.2 ABJ partition function

As a natural generalization of the ABJM case, we conjecture that the ABJ grand potential can be written in terms of the free energy of the refined topological string on local $\mathbb{P}^1\times\mathbb{P}^1$. For later convenience, we decompose the non-perturbative grand potential $J_{\rm np}(\mu)$ into the worldsheet instanton part $J_{\rm WS}(\mu)$, the D2-brane instanton part $J_{\rm D2}(\mu)$ and the mixed contributions $J_{\rm WS+D2}(\mu)$,

$$J_{\rm np}(\mu)=J_{\rm WS}(\mu)+J_{\rm D2}(\mu)+J_{\rm WS+D2}(\mu). \quad (4.23)$$

As in the ABJM case [2, 8, 17], we expect that the perturbative part plus the worldsheet instanton part is described by the un-refined topological string,

$$J_{\rm pert}(\mu)+J_{\rm WS}(\mu)=F_{\rm top}(T_1,T_2;g_s), \quad (4.24)$$

where $g_s=2/k$ in our definition (1.3). More explicitly, the worldsheet instanton part $J_{\rm WS}(\mu)$ is given by[8]

$$J_{\rm WS}(\mu)=\sum_{m=1}^{\infty}\sum_{d_{1,2},j_{L,R}}(-1)^{s_L+s_R+1}N^{d_1,d_2}_{j_L,j_R}\left[-\frac{s_R\sin\frac{4\pi m s_L}{k}}{m(2\sin\frac{2\pi m}{k})^2\sin\frac{4\pi m}{k}}e^{-m(d_1T_1+d_2T_2)}\right], \quad (4.25)$$

[8]Note that this expression is ill-defined for physical integer values of k. Hence this formula should be understood as the simple analytic continuation to unphysical values of k. As we will see in section 4.2.3, the divergences at integer values of k are actually apparent and canceled out if we take into account contributions from D2-brane instanton.

(d_1,d_2)	$\sum_{j_L,j_R} N^{d_1,d_2}_{j_L,j_R}(j_L,j_R)$
$(1,n), n\geq 0$	$(0,n+\frac{1}{2})$
$(2,2)$	$(\frac{1}{2},4)\oplus(0,\frac{7}{2})\oplus(0,\frac{5}{2})$
$(2,3)$	$(1,\frac{11}{2})\oplus(\frac{1}{2},5)\oplus(\frac{1}{2},4)\oplus 2(0,\frac{9}{2})\oplus(0,\frac{7}{2})\oplus(0,\frac{5}{2})$
$(2,4)$	$(\frac{3}{2},7)\oplus(1,\frac{13}{2})\oplus(1,\frac{11}{2})\oplus 2(\frac{1}{2},6)\oplus(\frac{1}{2},5)\oplus 2(0,\frac{11}{2})\oplus(\frac{1}{2},4)$ $\oplus 2(0,\frac{9}{2})\oplus(0,\frac{7}{2})\oplus(0,\frac{5}{2})$
$(3,3)$	$(2,\frac{15}{2})\oplus(\frac{3}{2},7)\oplus(\frac{3}{2},6)\oplus 3(1,\frac{13}{2})\oplus(\frac{1}{2},7)\oplus 2(1,\frac{11}{2})\oplus 3(\frac{1}{2},6)\oplus(1,\frac{9}{2})$ $\oplus 3(\frac{1}{2},5)\oplus 4(0,\frac{11}{2})\oplus 2(\frac{1}{2},4)\oplus 3(0,\frac{9}{2})\oplus(\frac{1}{2},3)\oplus 3(0,\frac{7}{2})\oplus(0,\frac{5}{2})\oplus(0,\frac{3}{2})$
$(2,5)$	$(2,\frac{17}{2})\oplus(\frac{3}{2},8)\oplus(\frac{3}{2},7)\oplus 2(1,\frac{15}{2})\oplus(1,\frac{13}{2})\oplus 2(\frac{1}{2},7)\oplus(1,\frac{11}{2})\oplus 2(\frac{1}{2},6)$ $\oplus 3(0,\frac{13}{2})\oplus(\frac{1}{2},5)\oplus 2(0,\frac{11}{2})\oplus(\frac{1}{2},4)\oplus 2(0,\frac{9}{2})\oplus(0,\frac{7}{2})\oplus(0,\frac{5}{2})$
$(3,4)$	$(3,\frac{19}{2})\oplus(\frac{5}{2},9)\oplus(\frac{5}{2},8)\oplus 3(2,\frac{17}{2})\oplus(\frac{3}{2},9)\oplus 2(2,\frac{15}{2})\oplus 4(\frac{3}{2},8)$ $\oplus(1,\frac{17}{2})\oplus(2,\frac{13}{2})\oplus 4(\frac{3}{2},7)\oplus 7(1,\frac{15}{2})\oplus 2(\frac{1}{2},8)\oplus(0,\frac{17}{2})\oplus 2(\frac{3}{2},6)$ $\oplus 6(1,\frac{13}{2})\oplus 7(\frac{1}{2},7)\oplus(0,\frac{15}{2})\oplus(\frac{3}{2},5)\oplus 5(1,\frac{11}{2})\oplus 8(\frac{1}{2},6)\oplus 7(0,\frac{13}{2})$ $\oplus 2(1,\frac{9}{2})\oplus 6(\frac{1}{2},5)\oplus 6(0,\frac{11}{2})\oplus(1,\frac{7}{2})\oplus 4(\frac{1}{2},4)\oplus 7(0,\frac{9}{2})\oplus 2(\frac{1}{2},3)$ $\oplus 4(0,\frac{7}{2})\oplus(\frac{1}{2},2)\oplus 3(0,\frac{5}{2})\oplus(0,\frac{3}{2})\oplus(0,\frac{1}{2})$

Table 1: BPS index $N^{d_1,d_2}_{j_L,j_R}$ of local $\mathbb{P}^1\times\mathbb{P}^1$ up to $d_1+d_2=7$.

where $N^{d_1,d_2}_{j_L,j_R}$ denotes the BPS index of local $\mathbb{P}^1\times\mathbb{P}^1$ partly listed[9] in table 1.

It is also expected that the D2-brane instanton part is captured by the Nekrasov-Shatashvili limit with the effective shift of the chemical potential,

$$J_{\rm pert}(\mu)+J_{\rm D2}(\mu)=J_{\rm pert}(\mu_{\rm eff})+\frac{1}{2\pi i}\frac{\partial}{\partial g_s}\left[g_s F_{\rm NS}\left(\frac{T_1^{\rm eff}}{g_s},\frac{T_2^{\rm eff}}{g_s};\frac{1}{g_s}\right)\right]. \tag{4.26}$$

In this expression, we treat $T^{\rm eff}_{1,2}$ and g_s as independent variables, namely

$$\frac{\partial T^{\rm eff}_{1,2}}{\partial g_s}=0. \tag{4.27}$$

As discussed in [2], the effective Kähler parameters are determined by the so-called quantum A-period, whose closed form is not known for the general value of g_s. However, for the physical case of integer k, we can write down the effective Kähler parameter $T^{\rm eff}_{1,2}$ in terms of a hypergeometric function (see appendix C for an explanation)

$$\begin{aligned}
\frac{T_I^{\rm eff}}{g_s}&=\frac{T_I}{g_s}-4e^{-\frac{T_I}{g_s}}\ {}_4F_3\left(1,1,\frac{3}{2},\frac{3}{2},2,2,2;16e^{-\frac{T_I}{g_s}}\right) \qquad \text{for} \quad g_s^{-1}\in\mathbb{Z}\\
\frac{2T_I^{\rm eff}}{g_s}&=\frac{2T_I}{g_s}-4e^{-\frac{2T_I}{g_s}}\ {}_4F_3\left(1,1,\frac{3}{2},\frac{3}{2},2,2,2;16e^{-\frac{2T_I}{g_s}}\right) \qquad \text{for} \quad g_s^{-1}\in\mathbb{Z}+\frac{1}{2},
\end{aligned} \tag{4.28}$$

which gives the effective chemical potential $\mu_{\rm eff}$ as

$$\mu_{\rm eff}=\begin{cases}\mu-2(-1)^{\frac{k}{2}-M}e^{-2\mu}\ {}_4F_3\left(1,1,\frac{3}{2},\frac{3}{2},2,2,2;(-1)^{\frac{k}{2}-M}16e^{-2\mu}\right) & \text{for } k:\text{even}\\ \mu+e^{-4\mu}\ {}_4F_3\left(1,1,\frac{3}{2},\frac{3}{2},2,2,2;-16e^{-4\mu}\right) & \text{for } k:\text{odd}\end{cases}. \tag{4.29}$$

[9] A part of the data is extracted from the table in section 5.5.2 of [63]. We are grateful to Can Kozçaz to tell us the other data of the BPS index, which is not listed in [63].

The second term of the right hand side in (4.26) takes the following explicit form

$$\begin{aligned}
&\frac{1}{2\pi i}\frac{\partial}{\partial g_s}\left[g_s F_{\rm NS}\left(\frac{T_1^{\rm eff}}{g_s},\frac{T_2^{\rm eff}}{g_s};\frac{1}{g_s}\right)\right] \\
&=\sum_{m=1}^{\infty}\sum_{d_{1,2},j_{L,R}}\frac{(-1)^{s_L+s_R+1}}{4m}N_{j_L,j_R}^{d_1,d_2} \\
&\left[\frac{\sin\frac{k\pi m s_L}{2}\sin\frac{k\pi m s_R}{2}}{\pi m\sin^3\frac{k\pi m}{2}}-\frac{k}{2}\frac{s_L\cos\frac{k\pi m s_L}{2}\sin\frac{k\pi m s_R}{2}+s_R\sin\frac{k\pi m s_L}{2}\cos\frac{k\pi m s_R}{2}}{\sin^3\frac{k\pi m}{2}}\right. \\
&\left.+\frac{k}{2\pi}\left(3\pi\cot\frac{k\pi m}{2}+d_1T_1^{\rm eff}+d_2T_2^{\rm eff}\right)\frac{\sin\frac{k\pi m s_L}{2}\sin\frac{k\pi m s_R}{2}}{\sin^3\frac{k\pi m}{2}}e^{-\frac{km}{2}(d_1T_1^{\rm eff}+d_2T_2^{\rm eff})}\right]. \qquad (4.30)
\end{aligned}$$

This shift of the chemical potential also plays an important role to describe the mixed contribution of the worldsheet instantons and D2-instantons by

$$J_{\rm WS}(\mu)+J_{\rm WS+D2}(\mu)=J_{\rm WS}(\mu_{\rm eff}). \qquad (4.31)$$

To summarize, we conjecture that the total ABJ grand potential is given by

$$\begin{aligned}
J_k^{(M)}(\mu)&=F_{\rm top}(T_1^{\rm eff},T_2^{\rm eff};g_s)+\frac{1}{2\pi i}\frac{\partial}{\partial g_s}\left[g_s F_{\rm NS}\left(\frac{T_1^{\rm eff}}{g_s},\frac{T_2^{\rm eff}}{g_s};\frac{1}{g_s}\right)\right] \\
&=\frac{C}{3}\mu_{\rm eff}^3+B\mu_{\rm eff}+A \\
&\quad+\sum_{m=1}^{\infty}\sum_{d_{1,2},j_{L,R}}(-1)^{s_L+s_R+1}N_{j_L,j_R}^{d_1,d_2} \\
&\quad\times\left[-\frac{s_R\sin\frac{4\pi m s_L}{k}}{m(2\sin\frac{2\pi m}{k})^2\sin\frac{4\pi m}{k}}e^{-m(d_1T_1^{\rm eff}+d_2T_2^{\rm eff})}\right. \\
&\quad+\frac{1}{4m}\left\{\frac{\sin\frac{k\pi m s_L}{2}\sin\frac{k\pi m s_R}{2}}{\pi m\sin^3\frac{k\pi m}{2}}-\frac{k}{2}\frac{s_L\cos\frac{k\pi m s_L}{2}\sin\frac{k\pi m s_R}{2}+s_R\sin\frac{k\pi m s_L}{2}\cos\frac{k\pi m s_R}{2}}{\sin^3\frac{k\pi m}{2}}\right. \\
&\quad\left.\left.+\frac{k}{2\pi}\left(3\pi\cot\frac{k\pi m}{2}+d_1T_1^{\rm eff}+d_2T_2^{\rm eff}\right)\frac{\sin\frac{k\pi m s_L}{2}\sin\frac{k\pi m s_R}{2}}{\sin^3\frac{k\pi m}{2}}\right\}e^{-\frac{km}{2}(d_1T_1^{\rm eff}+d_2T_2^{\rm eff})}\right]. \qquad (4.32)
\end{aligned}$$

We can rewrite this expression in the form of (3.10) and read off the coefficient $f_{l,m}(\mu)$ of the term $e^{-(2\ell+4m/k)\mu}$, as expected from the gravity side.

By exponentiating $J_k^{(M)}(\mu)$ in (3.10), we find the following expansion

$$e^{J_k^{(M)}(\mu)}=e^A\sum_{l,m=0}^{\infty}g_{l,m}(\mu)\exp\left[\frac{C}{3}\mu^3+\left(B-2l-\frac{4m}{k}\right)\mu\right], \qquad (4.33)$$

where $g_{l,m}(\mu)$ is a polynomial of μ explicitly determined by the coefficient $f_{l,m}(\mu)$. Then we come back to the canonical formalism by applying the integral transform (2.25) to (4.33),

$$\hat{Z}^{(N,N+M)}(k)=C^{-\frac{1}{3}}e^A\sum_{l,m=0}^{\infty}g_{l,m}\left(-\frac{\partial}{\partial N}\right)\mathrm{Ai}\left[C^{-\frac{1}{3}}\left(N-B+2l+\frac{4m}{k}\right)\right]. \qquad (4.34)$$

Since the Airy function satisfies the differential equation

$$\frac{d^2}{dz^2}\mathrm{Ai}[z] = z\mathrm{Ai}[z],$$

the canonical partition function given in (4.34) can be rewritten as a combination of the Airy function and its first derivative alone.

The expression (4.32) has apparent divergences at some values of the Chern-Simons level k, in particular at physical integer k. For example, for odd k such divergences occur when m in (4.32) is a multiple of k for the worldsheet instanton and even integer for D2-instanton, respectively. In the next subsubsection, we discuss that these divergences are actually canceled out and we compute the finite part of the grand potential.

4.2.3 Cancellation of divergence between instantons

Here we show the cancellation of apparent divergences in the grand potential (4.32) and compute its finite part.

Even Chern-Simons level

Let us first consider the even k case; we set $k = 2n_0$ with some integer n_0. By expanding[10] the m-th worldsheet instanton term with $m = n_0\ell$ ($\ell \in \mathbb{Z}$) around $k = 2n_0$, we find that the worldsheet instanton part (4.25) has poles at $k = 2n_0$:

$$\frac{(-1)^{s_L+s_R+(n+M)\ell d_-}}{4n_0\ell} s_R s_L N_{j_L,j_R}^{d_1,d_2} e^{-2ld_+\mu_{\mathrm{eff}}} \left[\frac{4n^2}{\pi^2\ell^2(k-2n_0)^2} + \frac{4n}{\pi^2\ell^2(k-2n_0)} + 1 + \frac{1}{\pi^2\ell^2} - \frac{2}{3}s_L^2 + \mathcal{O}(k-2n_0)\right], \tag{4.35}$$

where we have introduced the notation

$$d_\pm = d_1 \pm d_2. \tag{4.36}$$

The contribution (4.30) from F_{NS}, which gives the part of the D2-instanton correction, has also poles for arbitrary m:

$$\frac{(-1)^{(s_L+s_R+1)(mn_0+1)+mMd_-+d_-mn_0} e^{-2md_+\mu_{\mathrm{eff}}} s_L s_R N_{j_L,j_R}^{d_1,d_2}}{4m} \left[\frac{4n_0}{\pi^2m^2}\frac{1}{(k-2n_0)^2} + \frac{4}{\pi^2m^2}\frac{1}{k-2n_0} - \frac{n_0}{2} + \frac{n_0}{6}(s_L^2+s_R^2) - \frac{2d_+^2}{\pi^2n_0}\mu_{\mathrm{eff}}^2 - \frac{2d_+}{\pi^2mn_0}\mu_{\mathrm{eff}} + \frac{d_-^2}{2n_0}(n_0-M)^2 + \mathcal{O}(k-2n_0)\right]. \tag{4.37}$$

It is easy to see that these poles are canceled with each other if

$$(-1)^{s_L+s_R+1} = 1. \tag{4.38}$$

[10] This expansion should be handled with care. See the comments below and section 4.4 for detail.

It turns out that the non-zero values of the BPS index $N^{d_1,d_2}_{j_L,j_R}$ for local $\mathbb{P}^1\times\mathbb{P}^1$ appear only for the spins obeying the condition (4.38), and hence the poles are indeed canceled. After the pole cancellation, we find the finite part of the grand potential as[11]

$$\begin{aligned}
J(\mu) =& \frac{C}{3}\mu_{\rm eff}^3 + B\mu_{\rm eff} + A \\
&+ \sum_{m,2m/k\neq\mathbb{Z}}\sum_{d_{1,2},j_{L,R}} (-1)^{s_L+s_R+1} N^{d_1,d_2}_{j_L,j_R}\left[-\frac{e^{-2m\pi i d_-\left(\frac{1}{2}-\frac{M}{k}\right)} s_R \sin\frac{4\pi m s_L}{k}}{m(2\sin\frac{2\pi m}{k})^2\sin\frac{4\pi m}{k}} e^{-\frac{4m}{k}d_+\mu_{\rm eff}}\right] \\
&+ \sum_{m=1}^{\infty}\sum_{d_{1,2},j_{L,R}} \frac{s_L s_R}{2k} N^{d_1,d_2}_{j_L,j_R} \\
&\left[-\frac{2d_+^2}{\pi^2}\mu_{\rm eff}^2 {\rm Li}_1((-1)^{(\frac{k}{2}+M)d_+}e^{-2d_+\mu_{\rm eff}}) - \frac{2d_+}{\pi^2}\mu_{\rm eff}{\rm Li}_2((-1)^{(\frac{k}{2}+M)d_+}e^{-2d_+\mu_{\rm eff}})\right. \\
&-\frac{1}{\pi^2}{\rm Li}_3((-1)^{(\frac{k}{2}+M)d_+}e^{-2d_+\mu_{\rm eff}}) \\
&\left.+\left\{\frac{(16+k^2)s_L^2+k^2s_R^2}{24} - \frac{8+k^2}{8} + \frac{d_-^2}{2}\left(\frac{k}{2}-M\right)^2\right\}{\rm Li}_1((-1)^{(\frac{k}{2}+M)d_+}e^{-2d_+\mu_{\rm eff}})\right].
\end{aligned} \tag{4.39}$$

Some comments are in order here. First, when taking the limit $k\to 2n_0$ in our general expression (4.32), we should treat k and $T^{\rm eff}_{1,2}$ as independent variables since we have imposed the condition $\partial T^{\rm eff}_I/\partial g_s = 0$. This means that k in the expression of $T^{\rm eff}_{1,2}$ in (1.4) is set to $2n_0$ *before* taking the limit $k\to 2n_0$. Second, the resulting finite part (4.39) is invariant under the Seiberg-like duality $M\leftrightarrow k-M$ for even integer k. The same comments can be applied also to the odd k case.

Odd Chern-Simons level

When k is odd, there are also apparent divergences in the worldsheet instanton with $m=k\ell$ and the D2-instanton with $m=2\ell$ ($\ell\in\mathbb{Z}$). As in the even k case, similar calculation shows that these divergences are canceled again and the finite part of the grand potential is[12]

$$\begin{aligned}
J(\mu) =& \frac{C}{3}\mu_{\rm eff}^3 + B\mu_{\rm eff} + A \\
&+ \sum_{m,2m/k\neq\mathbb{Z}}\sum_{d_{1,2},j_{L,R}} (-1)^{s_L+s_R+1} N^{d_1,d_2}_{j_L,j_R}\left[-\frac{e^{-2m\pi i d_-\left(\frac{1}{2}-\frac{M}{k}\right)} s_R \sin\frac{4\pi m s_L}{k}}{m(2\sin\frac{2\pi m}{k})^2\sin\frac{4\pi m}{k}} e^{-\frac{4m}{k}d_+\mu_{\rm eff}}\right] \\
&+ \sum_{d_{1,2},j_{L,R}} \frac{s_L s_R}{4k} N^{d_1,d_2}_{j_L,j_R}\left[-\frac{2d_+^2}{\pi^2}\mu_{\rm eff}^2 {\rm Li}_1((-1)^{d_+}e^{-4d_+\mu_{\rm eff}}) - \frac{d_+}{\pi^2}\mu_{\rm eff}{\rm Li}_2((-1)^{d_+}e^{-4d_+\mu_{\rm eff}})\right. \\
&-\frac{1}{4\pi^2}{\rm Li}_3((-1)^{d_+}e^{-4d_+\mu_{\rm eff}}) \\
&\left.+\left\{\frac{(16+k^2)s_L^2+k^2s_R^2}{24} - \frac{8+k^2}{8} + \frac{d_-^2}{2}\left(\frac{k}{2}-M\right)^2\right\}{\rm Li}_1((-1)^{d_+}e^{-4d_+\mu_{\rm eff}})\right]
\end{aligned}$$

[11] We have used the definition of polylogarithm ${\rm Li}_s(z)=\sum_{p=1}^{\infty}\frac{z^p}{p^s}$.

[12] We have used $\sum_{p=1}^{\infty}\frac{x^{2p-1}}{2p-1} = {\rm Arctanh}x$.

$$+\frac{k}{8}\sum_{d_{1,2},j_{L,R}} N^{d_1,d_2}_{j_L,j_R} P_{d_+}\left(P_{s_R}s_R+P_{s_L}s_L\right)(-1)^{\frac{d_-}{2}+\frac{k}{2}(s_L+s_R+1)}\mathrm{Arctanh}(e^{-2d_+\mu_{\rm eff}}), \tag{4.40}$$

where P_l denotes the projection to even l,

$$P_l=\frac{1}{2}\Big(1+(-1)^l\Big). \tag{4.41}$$

4.3 Test of our proposal

Let us test our conjecture of the finite part (4.39) for even k and (4.40) for odd k, expected from the refined topological string. Since the odd number of D2-instanton does not appear in the finite part of the expected grand potential for odd k in (4.40), we can rewrite the expansion of the grand potential (4.34) as[13]

$$e^{J_k^{(M)}(\mu)}=e^A\sum_{m=0}^{\infty}\hat{g}_m(\mu)\exp\left[\frac{C}{3}\mu^3+\left(B-\frac{4m}{k}\right)\mu\right], \tag{4.42}$$

where $\hat{g}_m(\mu)$ is a polynomial of μ. Since we know the values of the BPS index $N^{d_1,d_2}_{j_L,j_R}$ up to $d_1+d_2=7$ from table 1, we can explicitly write down $\hat{g}_m(\mu)$ up to $m=7$. Using these coefficients $\hat{g}_m(\mu)$ $(m=0,\cdots,7)$, we define the putative partition function

$$\hat{Z}_{7-\rm inst}=e^A\sum_{m=0}^{7}\hat{g}_m\left(-\frac{\partial}{\partial N}\right)\mathrm{Ai}\left[C^{-1/3}\left(N-B+\frac{4m}{k}\right)\right]. \tag{4.43}$$

If our conjecture is correct, then this quantity should be the canonical partition function including the instanton effects up to the order of 7-th worldsheet instanton correction. Hence we expect that the difference between the exact partition function $\hat{Z}_k^{(M)}$ and $\hat{Z}_{7-\rm inst}$ behaves as

$$\frac{\hat{Z}_k^{(M)}-\hat{Z}_{7-\rm inst}}{\hat{Z}_{\rm pert}}=\frac{\sum_{m=8}^{\infty}\hat{g}_m\left(-\frac{\partial}{\partial N}\right)\mathrm{Ai}\Big[C^{-1/3}\left(N-B+\frac{4m}{k}\right)\Big]}{\mathrm{Ai}\Big[C^{-1/3}\left(N-B\right)\Big]}=\mathcal{O}(N^{\frac{\alpha}{2}}e^{-16\pi\sqrt{\frac{2N}{k}}}), \tag{4.44}$$

in the large N regime. Here α is some non-negative integer depending[14] on k and M. In fig. 4, we plot the quantity

$$e^{14\pi\sqrt{2N}k}\frac{\left|\hat{Z}_k^{(M)}-\hat{Z}_{7-\rm inst}\right|}{\hat{Z}_{\rm pert}}, \tag{4.45}$$

against $2\pi\sqrt{2N/k}$ both in semi-log scale (Left) and log-log scale (Right) for various values of (k,M). From fig. 4 one can see that this quantity (4.45) is extremely small, and this already

[13]Note that this expression of course includes also D2-instanton effects since $k\ell/2$-th worldsheet instanton actually contributes by the same weight as ℓ-th D2-instanton.

[14]For example, if (k,M) is $(2,1)$, the formula (4.39) implies $\alpha=16$. This is because the grand partition function contains a term with $\mathcal{O}(\mu^{16}e^{-16\mu})$ coming from the μ^2-terms in the first D2-instanton correction of the grand potential.



Figure 4: The quantity $e^{14\pi\sqrt{2N}k}|\hat{Z}_k^{(M)} - \hat{Z}_{7-\text{inst}}|/\hat{Z}_{\text{pert}}$ is plotted to $2\pi\sqrt{2N/k}$ both in semi-log scale (Left) and log-log scale (Right) for $(k, M) = (2,1), (3,1), (4,1), (4,2), (6,1), (6,2)$ and $(6,3)$. The blue circle, purple square and yellow diamond symbols show the cases for $M = 1$, $M = 2$ and $M = 3$, respectively.

gives strong evidence for our conjecture. We can argue more precisely as follows. If $\hat{Z}_{7-\text{inst}}$ properly contains the instanton effects, then this quantity (4.45) should be exponentially suppressed by $e^{-2\pi\sqrt{\frac{2N}{k}}}$ in the large N regime. On the other hand, if $\hat{Z}_{7-\text{inst}}$ had wrong information at 6-th or lower order of worldsheet instanton effects, we could observe an exponential blow-up of this quantity. The semi-log plots in fig. 4 (Left) show that such exponential blow-up does not occur. We also see from the log-log plots in fig. 4 (Right) that the data points give a power law behavior in not so large N region. This behavior is consistent with our expectation (4.44) since the data should show such a power law behavior before going to exponentially suppressed region. Thus we conclude that the grand potential (4.32) coming from the refined topological string is correct at least up to the order of 6-th worldsheet instanton corrections. In particular the case of $(k, M) = (4, 1)$, plotted in the second bottom of fig. 4 (Left), clearly exhibits the exponentially suppressed behavior. This strongly supports our expectation (4.44).

To summarize, all cases we studied strongly support our conjecture that the ABJ grand potential is given by the general expression (1.2) in terms of the refined topological string on local $\mathbb{P}^1 \times \mathbb{P}^1$, with an appropriate identification of the Kähler parameters (1.4).

4.4 Comments on Matsumoto-Moriyama proposal

Recently an apparently similar form of the grand potential was proposed in [30] by a different approach, which uses a relation between the ABJ partition function and the half-BPS Wilson loops of the ABJM theory. The work [30] have calculated the ABJ partition function exactly or numerically with high precision for $(k, M, N_{\max}) = (2, 1, 4)$, $(3, 1, 3)$, $(4, 1, 2)$, $(4, 2, 2)$, $(6, 1, 3)$, $(6, 2, 2)$ and $(6, 3, 2)$. These values are totally consistent with our exact values listed in appendix D. However, there is an important difference of physical interpretation between [30] and ours.

This difference comes from an ambiguity for taking the limit to a physical Chern-Simons level in the apparently divergent grand potential (4.32). Suppose that we take the limit $k \to k_0$ ($k_0 \in \mathbb{Z}$) in (4.32). In section 4.2.3, we have expanded (4.32) around $k = k_0$ after fixing $k = k_0$ in the effective Kähler parameters T_I^{eff} since we have imposed $\partial T_I^{\text{eff}}/\partial g_s = 0$, while the authors in [30] expanded the total expression (4.32) around $k = k_0$. This difference of taking the limit affects the finite part[15] of the grand potential after the cancellation of divergence. Since the latter prescription does not agree with the exact data, the authors in [30] added an extra new term, whose origin is unclear from the topological string perspective, in such a way that the finite part agrees with their exact and numerical data. It turns out that the finite part of our grand potential, corresponding to (4.39) and (4.40), agrees with the one given in Figure 4 of [30]. However, our grand potential before cancellation (4.32) is different from the proposal in [30] with an extra term added. We believe that our prescription is more natural since everything can be completely explained by the refined topological string.

[15] Fortunately or unfortunately, this difference vanishes for $M = 0$. Therefore this ambiguity of how to take the limit is irrelevant for the ABJM case [2].

5. Discussions

In this paper we have studied the partition function of the ABJ theory on S^3. By using the Fermi gas formalism and Tracy-Widom's lemma [47], we have exactly computed the ABJ partition function for $(k, M, N_{\max}) = (2, 1, 65)$, $(3, 1, 31)$, $(4, 1, 62)$, $(4, 2, 29)$, $(6, 1, 23)$, $(6, 2, 21)$ and $(6, 3, 22)$. These exact data enable us to test the relation between the ABJ partition function and the free energy of the refined topological string on local $\mathbb{P}^1 \times \mathbb{P}^1$ with non-diagonal Kähler parameters. We conclude that the ABJ partition function is completely determined by the refined topological string including full series of the membrane instanton effects in the M-theory dual.

Although here we have focused on the usual AdS/CFT correspondence between the ABJ theory and the M-theory on $AdS_4 \times S^7/\mathbb{Z}_k$, it is also interesting to study the higher spin limit [28, 29], corresponding to the regime where $M, k \gg 1$ with M/k and N kept fixed. It would also be interesting to study the eigenvalue problem of the Hamiltonian (3.6) of the ABJ Fermi gas, as in the case of the ABJM theory [20]. We leave this for future work.

Throughout many previous works about the ABJM theory and this work, we have seen that the (refined) topological string is a very powerful tool to determine the structures of the partition function and the BPS Wilson loops in the ABJ(M) theory. However, if we consider more general M2-brane theory beyond[16] the ABJ(M) theory, the correspondence to the topological string is rather obscure, and it might be the case that there is no such relation in general. Therefore even if we can exactly compute the partition function for the case other than the ABJ(M) theory for some discrete values of parameters, we do not have a good guiding principle to extend the non-perturbative effects to more general parameters. Although this problem can be overcome for the "orbifold ABJ(M) theory" [46] due to a relation to the ABJ(M) theory, we need to develop a new approach to study more general M2-brane theories. It is very interesting to study the instanton effects in other M2-brane theories having AdS_4 dual.

Acknowledgment

The authors would like to thank Yasuyuki Hatsuda, Johan Källén, Hiroaki Kanno, Can Kozçaz, Sanefumi Moriyama, Kazuhiro Sakai, Ashoke Sen and Masato Taki for valuable discussions. We would also like to thank Satoru Odake for allowing us to use computers in the theory group, Shinshu University, for the computation of the exact partition functions. M.H. is grateful to KEK Theory Center and Kavli IPMU for warm hospitality. The work of K.O. is supported in part by JSPS Grant-in-Aid for Young Scientists (B) 23740178.

A. Algorithm for exact computation of ABJ grand partition function

In this appendix, we show that the grand canonical partition function of the ABJ theory is determined by a series of functions $\phi_l^+(y)$ which can be computed recursively by the

[16] For previous studies, see e.g. [64, 65, 66, 67, 68, 69] in the M-theory limit, [70, 71, 72, 73] in the 't Hooft limit and [13, 55, 74, 75, 46] in the Fermi gas approach.

formula (2.27). Our algorithm is essentially a simple generalization of the ABJM case [16, 17].

As discussed in section 2, the ABJ grand partition function is given by

$$\Xi_k^{(M)}(z) = \exp\left[-\sum_{n=1}^{\infty}\frac{(-1)^n}{n}\mathrm{Tr}\rho^n\right], \tag{A.1}$$

with the density matrix (2.8) . Here we define multiplication and trace for certain matrices ρ_1, ρ_2 as

$$\rho_1\rho_2(x,y) = \int_{-\infty}^{\infty}\frac{dq}{2\pi k}\rho_1(x,q)\rho_2(q,y), \quad \mathrm{Tr}\rho_1 = \int_{-\infty}^{\infty}\frac{dq}{2\pi k}\rho_1(q,q), \tag{A.2}$$

respectively. Then we decompose the density matrix into the even and odd parts:

$$\rho(x,y) = \rho_+(x,y) + \rho_-(x,y), \quad \rho_\pm(x,y) = \frac{\rho(x,y)\pm\rho(x,-y)}{2}. \tag{A.3}$$

In terms of $\rho_\pm$, we rewrite the grand partition function as

$$\Xi_k^{(M)}(z) = \Xi_+(z)\Xi_-(z), \quad \Xi_\pm(z) = \mathrm{Det}(1+z\rho_\pm). \tag{A.4}$$

Note that $\rho_\pm$ takes the form of

$$\rho_\pm(x,y) = \frac{E_\pm(x)E_\pm(y)}{\cosh\frac{x}{k}+\cosh\frac{y}{k}}, \quad E_\pm(x) = \frac{\sqrt{V(x)}\left(e^{\frac{x}{2k}}\pm e^{\frac{x}{2k}}\right)}{2}. \tag{A.5}$$

Therefore, both $\rho_\pm$ have the form of

$$\frac{E(x)E(y)}{M(x)+M(y)},$$

where the Tracy-Widom's lemma [47] is applicable. Then the power of $\rho_\pm$ is determined by a series of functions $\phi_l^\pm(y)$

$$\begin{aligned}
\rho_\pm^{2n+1}(x,y) &= \frac{E_\pm(x)E_\pm(y)}{\cosh\frac{x}{k}+\cosh\frac{y}{k}}\sum_{l=0}^{2n}(-1)^l\phi_l^\pm(x)\phi_{2n-l}^\pm(y),\\
\rho_\pm^{2n}(x,y) &= \frac{E_\pm(x)E_\pm(y)}{\cosh\frac{x}{k}-\cosh\frac{y}{k}}\sum_{l=0}^{2n-1}(-1)^l\phi_l^\pm(x)\phi_{2n-1-l}^\pm(y),\\
\phi_l^\pm(x) &= \frac{1}{E_\pm(x)}\int_{-\infty}^{\infty}\frac{dq}{2\pi k}\rho_\pm^l(x,q)E_\pm(q), \quad \phi_0^\pm(x) = 1.
\end{aligned} \tag{A.6}$$

At this stage, it seems that we should find the two series of the functions $\phi_l^+(y)$ and $\phi_l^-(y)$ to obtain the grand potential. However, similar to the ABJM case [17], one can show the following convenient identity,

$$\frac{\Xi_-(z)}{\Xi_+(-z)} = \sum_{l=0}^{\infty}\phi_l^+(0)z^l, \tag{A.7}$$

which implies that we need information of only $\phi_l^+(y)$. Plugging this into (A.4), we arrive at (2.26). In the rest of this section, we present a concrete algorithm to compute $\phi_l^+(y)$ and $\mathrm{Tr}\rho_+^{2n}$ exactly.

A.1 Algorithm for even Chern-Simons level

By changing variables as $u = e^{\frac{y}{k}}, v = e^{\frac{y'}{k}}$, the recursion equation (2.27) becomes

$$\phi^+_{l+1}(u) = \frac{1}{2\pi}\frac{u}{u+1}\int_0^\infty dv \frac{v^{\frac{k}{2}-1}}{u+v}\frac{v+1}{v^k+(-1)^M}\left[\prod_{s=-\frac{M-1}{2}}^{\frac{M-1}{2}}\frac{v-e^{-\frac{2\pi s}{k}}}{v+e^{-\frac{2\pi s}{k}}}\right]\phi^+_l(v). \tag{A.8}$$

If we assume the form of $\phi^+_l(u)$ as

$$\phi^+_l(u) = \sum_{j=0}^{l} A^{(j)}_l(u)(\log u)^j, \tag{A.9}$$

with rational functions $A^{(j)}_l$, we find[17]

$$\begin{aligned}\phi^+_{l+1}(u) = &-\frac{1}{2\pi}\frac{u}{u+1}\sum_{j=0}^{l}\frac{(2\pi i)^{j+1}}{j+1} \\ &\sum_{\text{poles}}\text{Res}_v\left[A^{(j)}_l(v)B_{j+1}\left(\frac{\log v}{2\pi i}\right)\frac{v^{\frac{k}{2}-1}}{u+v}\frac{v+1}{v^k+(-1)^M}\prod_{s=-\frac{M-1}{2}}^{\frac{M-1}{2}}\frac{v-e^{-\frac{2\pi i s}{k}}}{v+e^{-\frac{2\pi i s}{k}}}\right].\end{aligned} \tag{A.10}$$

The symbol $\sum_{\text{poles}}$ indicates the summation over all the poles of the integrand. From $\phi^+_l(u)$, we can find $\rho^{2n}_+(u,u)$ as

$$\rho^{2n}_+(u,u) = \frac{1}{2}\frac{u^{\frac{k}{2}+1}(u+1)}{(u-1)(u^k+(-1)^M)}\prod_{s=-\frac{M-1}{2}}^{\frac{M-1}{2}}\left[\frac{u-e^{-\frac{2\pi i s}{k}}}{u+e^{-\frac{2\pi i s}{k}}}\right]\sum_{l=0}^{2n-1}(-1)^l\frac{d\phi^+_l(u)}{du}\phi^+_{2n-l-1}(u). \tag{A.11}$$

By expanding $\rho^{2n}_+(u,u)/u$ in terms of the power of $\log u$

$$\frac{1}{u}\rho^{2n}_+(u,u) = \sum_{j=0}^{2n-1} R^{(j)}_n(u)(\log u)^j, \tag{A.12}$$

where $R^{(j)}_n(u)$ is some rational function of u, then the trace of ρ^{2n}_+ can be obtained as a sum of residues

$$\text{Tr}\,\rho^{2n}_+ = -\frac{1}{2\pi}\sum_{j=0}^{2n-1}\frac{(2\pi i)^{j+1}}{j+1}\sum_{\text{poles}}\text{Res}_u\left[R^{(j)}_n(u)B_{j+1}\left(\frac{\log u}{2\pi i}\right)\right]. \tag{A.13}$$

[17] Here we have used an integral formula noted in [16],

$$\int_0^\infty dv C(v)\log^j v = -\frac{(2\pi i)^j}{j+1}\oint_\gamma dv C(v)B_{j+1}\left(\frac{\log v}{2\pi i}\right),$$

where $C(v)$ is a rational function and B_{j+1} is the Bernoulli polynomial. Choosing the branch cut of $\log v$ as the positive real axis, the integral contour γ goes from $+\infty$ to 0 infinitesimally below the branch cut and then to $+\infty$ infinitesimally above the branch cut.

A.2 Algorithm for odd Chern-Simons level

When k is odd, by introducing the variables $u = e^{y/2k}, v = e^{y'/2k}$, we can rewrite the recursion relation (2.27) as

$$\phi^+_{l+1}(u) = \frac{1}{2\pi}\frac{u^2}{u^2+1}\int_0^\infty dv \frac{v^{k-1}}{u^2+v^2}\frac{v^2+1}{v^{2k}+(-1)^M}\left[\prod_{s=-\frac{M-1}{2}}^{\frac{M-1}{2}}\frac{v^2-e^{-\frac{2\pi i s}{k}}}{v^2+e^{-\frac{2\pi i s}{k}}}\right]\phi^+_l(v). \tag{A.14}$$

By expanding ϕ^+_{l+1} as (A.9), we find

$$\phi^+_{l+1}(u) = -\frac{1}{\pi}\frac{u^2}{u^2+1}\sum_{j=0}^{l}\frac{(2\pi i)^{j+1}}{j+1} \sum_{\text{poles}}\text{Res}_v\left[A^{(j)}_l(v)B_{j+1}\left(\frac{\log v}{2\pi i}\right)\frac{v^{k-1}}{u^2+v^2}\frac{v^2+1}{v^{2k}+(-1)^M}\prod_{s=-\frac{M-1}{2}}^{\frac{M-1}{2}}\frac{v^2-e^{-\frac{2\pi i s}{k}}}{v^2+e^{-\frac{2\pi i s}{k}}}\right]. \tag{A.15}$$

Then $\rho^{2n}_+(u,u)$ is rewritten as

$$\rho^{2n}_+(u,u) = \frac{1}{4}\frac{u^{k+1}(u^2+1)}{(u^2-1)(u^{2k}+(-1)^M)}\prod_{s=-\frac{M-1}{2}}^{\frac{M-1}{2}}\left[\frac{u^2-e^{-\frac{2\pi i s}{k}}}{u^2+e^{-\frac{2\pi i s}{k}}}\right]\sum_{l=0}^{2n-1}(-1)^l\frac{d\phi^+_l(u)}{du}\phi^+_{2n-l-1}(u). \tag{A.16}$$

In a similar way as the even k case, by expanding $\rho^{2n}_+(u,u)/u$ in terms of the power of $\log u$

$$\frac{1}{u}\rho^{2n}_+(u,u) = \sum_{j=0}^{2n-1}R^{(j)}_n(u)(\log u)^j, \tag{A.17}$$

the trace of ρ^{2n}_+ can be obtained as a sum of residues

$$\text{Tr}\,\rho^{2n}_+ = -\frac{1}{\pi}\sum_{j=0}^{2n-1}\frac{(2\pi i)^{j+1}}{j+1}\sum_{\text{poles}}\text{Res}_u\left[R^{(j)}_n(u)B_{j+1}\left(\frac{\log u}{2\pi i}\right)\right]. \tag{A.18}$$

B. π^{-N} term of $\widehat{Z}^{(N,N+M)}(k)$

In this appendix, we will briefly discuss the structure of the highest transcendental term of the partition functions of the ABJ theory listed in appendix D, and the partition functions of the ABJM theory in [17]. As mentioned in [17] for the $k = 1$ ABJM theory, the coefficient of the highest transcendental term π^{-N} in the canonical partition function can be expressed in terms of the Hermite polynomial using the relation

$$e^{2xz-z^2} = \sum_{n=0}^{\infty}\frac{H_n(x)}{n!}z^n. \tag{B.1}$$

We will see that the partition functions of ABJ theory also have a similar structure.

B.1 $(k, M) = (2, 1)$

The partition function of $(k, M) = (2, 1)$ case has the structure

$$\widehat{Z}^{(N,N+1)}(2) = \sum_{m=0}^{N} \frac{c_m^{(N)}}{\pi^m}. \tag{B.2}$$

Let us consider the highest transcendental term π^{-N} in $\widehat{Z}^{(N,N+1)}(2)$. We observe that such highest term of $\phi_l(u=1)$ comes from the $(\log u)^l$ term in the expansion (A.9)

$$\begin{aligned}
\phi_l^+(u) &= \varepsilon_l \frac{\phi_1^+(u)^l}{l!} + \cdots, \qquad \varepsilon_l = \sqrt{2}\sin\left(\frac{\pi(2l+1)}{4}\right) = \pm 1, \\
\phi_l^+(1)_{\text{highest}} &= \varepsilon_l \frac{\phi_1^+(1)^l}{l!} = \frac{\varepsilon_l}{(4\pi)^l l!},
\end{aligned} \tag{B.3}$$

where

$$\phi_1^+(u) = \frac{u}{2\pi(u^2-1)} \log u. \tag{B.4}$$

$\operatorname{Tr} \rho_+^{2m}$ with $m \geq 2$ is always less transcendental compared to π^{-2m}, and the only highest transcendental term comes from $\operatorname{Tr} \rho_+^2 = \frac{1}{32\pi^2}$. Thus we find that the generating function of the highest term π^{-N} is given by

$$\begin{aligned}
\sum_{N=0}^{\infty} z^N \frac{c_N^{(N)}}{\pi^N} &= e^{-\operatorname{Tr}\rho_+^2 z^2} \left(\sum_{l=0}^{\infty} \phi_l^+(1)_{\text{highest}} z^l \right) \\
&= e^{-\frac{z^2}{32\pi^2}} \left(\cos\frac{z}{4\pi} + \sin\frac{z}{4\pi} \right) \\
&= 1 + \frac{1}{4\pi} z - \frac{1}{16\pi^2} z^2 - \frac{1}{96\pi^3} z^3 + \frac{5}{3072\pi^4} z^4 + \cdots.
\end{aligned} \tag{B.5}$$

We further find

$$\sum_{m=1}^{\infty} \frac{\operatorname{Tr} \rho_+^{2m}}{m} z^{2m} = \frac{z^2}{2^5\pi^2} + \frac{z^4}{2^9 3\pi^2} + \frac{z^6}{2^{10}3^3 5\pi^2} + \frac{z^6}{2^{16}\pi^2} - \frac{z^4}{2^{15}} \left(1 + \cos\frac{z}{\pi}\right) + \cdots. \tag{B.6}$$

B.2 $(k, M) = (4, 1)$

In a similar manner, the highest transcendental term for the $(k, M) = (4, 1)$ case is found to be

$$\begin{aligned}
\sum_{N=0}^{\infty} z^N \frac{c_N^{(N)}}{\pi^N} &= e^{-\operatorname{Tr}(\rho_+^2)_{\text{highest}} z^2} \left(\sum_{l=0}^{\infty} \phi_l^+(1)_{\text{highest}} z^l \right) \\
&= \exp\left(\frac{z^2}{64\pi^2} - \frac{z}{8\pi} \right) \\
&= 1 - \frac{1}{8\pi} z + \frac{3}{128\pi^2} z^2 - \frac{7}{3072\pi^3} z^3 + \frac{25}{98304\pi^4} z^4 + \cdots.
\end{aligned} \tag{B.7}$$

The next-to-highest term is

$$\sum_{N=1}^{\infty} z^N \frac{c_{N-1}^{(N)}}{\pi^{N-1}} = \frac{z}{16} \exp\left(\frac{z^2}{64\pi^2} + \frac{3z}{8\pi}\right). \tag{B.8}$$

Therefore, we find

$$\Xi(z) = e^{\frac{z^2}{64\pi^2} - \frac{z}{8\pi}} \left[1 + \frac{z}{16} e^{\frac{z}{2\pi}} + \cdots\right]. \tag{B.9}$$

B.3 Summary of the Higher Transcendental Part of Grand Partition Functions

By inspecting the exact partition functions, we find a similar structure for other cases. Here we summarize the higher transcendental part of grand partition functions $\Xi_k^{(M)}(z)$ for various k and M,

$$\begin{aligned}
\Xi_1^{(0)}(z) &= \exp\left[\frac{z^4}{2^7\pi^2} + \frac{z^2}{2^4\pi}\cdots\right] \cdot \left[1 + \frac{z}{4} e^{-\frac{z^2}{4\pi}} + \frac{z^3}{64} e^{\frac{z^2}{4\pi}} \cdots\right] \\
\Xi_3^{(0)}(z) &= \exp\left[\frac{z^4}{3\cdot 2^7\pi^2} - \frac{z^2}{2^4\pi} + \cdots\right] \cdot \left[1 + \frac{z}{12} e^{\frac{z^2}{12\pi}} + \cdots\right] \\
\Xi_2^{(0)}(z) &= \exp\left[\frac{z^2}{2\cdot 2^4\pi^2} - \frac{z^4}{2^9\cdot 3\pi^2} + \frac{z^6}{2^{10}3^3 5\pi^2} + \frac{z^6}{2^{16}\pi^2} + \frac{z^4}{2^{14}} \cosh^2\frac{z}{2\pi} + \cdots\right] \\
&\quad \cdot \left[1 + \frac{z}{8}\cosh\frac{z}{2\pi} + \cdots\right] \\
\Xi_2^{(1)}(z) &= \exp\left[-\frac{z^2}{2\cdot 2^4\pi^2} - \frac{z^4}{2^9\cdot 3\pi^2} - \frac{z^6}{2^{10}3^3 5\pi^2} - \frac{z^6}{2^{16}\pi^2} + \frac{z^4}{2^{14}} \cos^2\frac{z}{2\pi} + \cdots\right] \\
&\quad \cdot \left[\cos\frac{z}{4\pi} + \sin\frac{z}{4\pi}\cdots\right] \\
\Xi_4^{(0)}(z) &= \exp\left[-\frac{z^2}{4\cdot 2^4\pi^2} + \cdots\right] \cdot \left[1 + \frac{z}{16} - \frac{z^2}{128}\sin\frac{z}{2\pi} + \cdots\right] \\
\Xi_4^{(1)}(z) &= \exp\left[\frac{z^2}{4\cdot 2^4\pi^2} - \frac{z}{8\pi} + \cdots\right] \cdot \left[1 + \frac{z}{16} e^{\frac{z}{2\pi}} + \cdots\right] \\
\Xi_6^{(0)}(z) &= \exp\left[\frac{z^2}{6\cdot 2^4\pi^2} + \cdots\right] \cdot \left[1 + \frac{z}{24}\cosh\frac{z}{6\pi} + \frac{\sqrt{3}z^2}{2\cdot 3^4}\sinh\frac{z}{6\pi} + \cdots\right] \\
\Xi_6^{(1)}(z) &= \exp\left[-\frac{z^2}{6\cdot 2^4\pi^2} + \cdots\right] \cdot \left[\cos\frac{z}{12\pi} + \sin\frac{z}{12\pi} + \cdots\right].
\end{aligned} \tag{B.10}$$

We do not have a clear understanding of the origin of this structure. We leave this for a future work.

C. Picard-Fuchs equation and effective Kähler parameters

In this appendix, we will explain that the effective Kähler parameters for integer k are given by (4.28), which are essentially equal to the classical periods of diagonal local $\mathbb{P}^1 \times \mathbb{P}^1$.

The Picard-Fuchs equation of local $\mathbb{P}^1 \times \mathbb{P}^1$ for the classical periods Π is

$$\theta_I^2 \Pi = z_I(2\theta_1 + 2\theta_2)(2\theta_1 + 2\theta_2 + 1)\Pi, \qquad (I = 1, 2) \tag{C.1}$$

with

$$\theta_I = z_I \frac{\partial}{\partial z_I}. \tag{C.2}$$

Note that θ_I measures the exponent of z_I^n

$$\theta_I z_I^n = n z_I^n. \tag{C.3}$$

The classical A-period is a solution of the Picard-Fuchs equation (C.1)

$$\Pi_{A_I} = \log z_I + 2 \sum_{k,l \geq 0} \frac{(2k+2l-1)!}{(k!l!)^2} z_1^k z_2^l. \tag{C.4}$$

For the diagonal case $z_1 = z_2 = z$, this reduces to

$$\Pi_A = \log z + 4z \sum_{n=0}^{\infty} \frac{(2n+1)!^2}{(n+1)!^3} \frac{z^n}{n!} = \log z + 4z {}_4F_3 \left(1, 1, \frac{3}{2}, \frac{3}{2}, 2, 2, 2; 16z\right), \tag{C.5}$$

where we have used the identity

$$\sum_{m=0}^{n} \binom{n}{m}^2 = \binom{2n}{n}. \tag{C.6}$$

As discussed in [17], the effective Kähler parameter is given by the quantum A-period. For even k, the quantum parameter q is equal to the classical value $q = e^{i\pi k} = 1$, and the instanton factors for the D2-brane become "diagonal"

$$z = e^{-T_{1,2}/g_s} = (-1)^{\frac{k}{2}-M} e^{-2\mu}, \tag{C.7}$$

although the worldsheet instanton factors are non-diagonal $e^{-T_1} \neq e^{-T_2}$. Therefore, the quantum A-period is reduced to the classical period (C.5). In [19], it is conjectured for the ABJM case that the quantum period for the odd k case, corresponding to $q = -1$, is also expressed in terms of the hypergeometric function, obtained by simply replacing $z \to z^2$ in (C.5). We expect that this is also the case for the ABJ theory.

D. Exact results on ABJ partition function

In this section, we present exact results of the ABJ partition function for various k, M and N. Although we have indeed obtained the exact values for $(k, M, N_{\max}) = (2, 1, 65)$, $(3, 1, 31)$, $(4, 1, 62)$, $(4, 2, 29)$, $(6, 1, 23)$, $(6, 2, 21)$ and $(6, 3, 22)$, we list these values up to $N = 20$ at most. This is because the expressions for larger N are too long. The other values are available upon request to the present authors. Our values are consistent with the result obtained in the work [30] by a different approach, which has calculated the ABJ partition function exactly or numerically with high precision for $(k, M, N_{\max}) = (2, 1, 4)$, $(3, 1, 3)$, $(4, 1, 2)$, $(4, 2, 2)$, $(6, 1, 3)$, $(6, 2, 2)$ and $(6, 3, 2)$.

$$\hat{Z}^{(1,2)}(2)=\frac{1}{4\pi},\quad \hat{Z}^{(2,3)}(2)=\frac{1}{128}-\frac{1}{16\pi^2},\quad \hat{Z}^{(3,4)}(2)=\frac{5\pi^2-48}{4608\pi^3},\quad \hat{Z}^{(4,5)}(2)=\frac{9}{32768}+\frac{5}{3072\pi^4}-\frac{53}{18432\pi^2},\quad \hat{Z}^{(5,6)}(2)=\frac{6240-800\pi^2+17\pi^4}{29491200\pi^5},$$

$$\hat{Z}^{(6,7)}(2)=\frac{-218880+1413600\pi^2-1160264\pi^4+103275\pi^6}{8493465600\pi^6},\quad \hat{Z}^{(7,8)}(2)=\frac{-4677120-8631840\pi^2+14206864\pi^4-1345977\pi^6}{1664719257600\pi^7},$$

$$\hat{Z}^{(8,9)}(2)=\frac{61608960-1051438080\pi^2+2363612608\pi^4-1477376224\pi^6+126511875\pi^8}{213084064972800\pi^8},\quad \hat{Z}^{(9,10)}(2)=\frac{633830400+6140897280\pi^2-22473501120\pi^4+16465544384\pi^6-1444050207\pi^8}{23013079017062400\pi^9},$$

$$\hat{Z}^{(10,11)}(2)=\frac{-45945446400+1789603200000\pi^2-8040102013440\pi^4+12612559256000\pi^6-6759848237256\pi^8+563610403125\pi^{10}}{1841046321364992000\pi^{10}},$$

$$\hat{Z}^{(11,12)}(2)=-\frac{1899826790400+58020665472000\pi^2-380149120834560\pi^4+713711875201600\pi^6-419001890609104\pi^8+35516095421175\pi^{10}}{8910664195406561280000\pi^{11}},$$

$$\hat{Z}^{(12,13)}(2)=\frac{1}{513254257655417929728000\pi^{12}}\Big[8951070965760-730576220774400\pi^2+55891102669839360\pi^4-16567100527296000\pi^6+2125955978162498\pi^8$$
$$-102685378935079440\pi^{10}+8388213629709375\pi^{12}\Big],$$

$$\hat{Z}^{(13,14)}(2)=\frac{1}{34695987817506252049612800000\pi^{13}}\Big[4720114977177600+362547424176537600\pi^2-3751225073125770240\pi^4+13042072681841448960\pi^6$$
$$-19012094772801329056\pi^8+9993495094908215904\pi^{10}-8305493191234706\pi^{12}\Big],$$

$$\hat{Z}^{(14,15)}(2)=\frac{1}{5440330889784980321379287040000\pi^{14}}\Big[-555727897460736000+89573494835323699200\pi^2-10556361504673560576\pi^4+513148883682802278912\pi^6$$
$$-12078328057432325328640\pi^8+1353783170736361458620\pi^{10}-60518928035620436410\pi^{12}+4862395794733633406\pi^{14}\Big],$$

$$\hat{Z}^{(15,16)}(2)=-\frac{1}{4896297800806482289241358336000000\pi^{15}}\Big[360901945277153280000+61045839496715673600000\pi^2-9206750935311831982080\pi^4$$
$$+507831737592928484736000\pi^6-13440434769822663713510\pi^8+1708199914796799315018400\pi^{10}-84103881813497711786558\pi^{12}+6902417670115186756687\pi^{14}\Big],$$

$$\hat{Z}^{(16,17)}(2)=\frac{1}{125345223700645946604578773401600000\pi^{16}}\Big[644515885825523712000-195903374317541130240000\pi^2+331778842551153816698880\pi^4-24242575894767562235904000\pi^6$$
$$+9168657937760942429512704\pi^8-1846214972763849411616256\pi^{10}+187135567910967396538249344\pi^{12}-7870540152158521604405280\pi^{14}+623602260674588484351562\pi^{16}\Big],$$

$$\hat{Z}^{(17,18)}(2)=\frac{1}{14489907859794671427489306205224960000\pi^{17}}\Big[502229017051881799680+17114872531857226334208000\pi^2-35582518059145524674887680\pi^4$$
$$+28386738978977086356160512\pi^6-11715836542518334641324349440\pi^8+26594007390595358524134338560\pi^{10}$$
$$-31088486157208526910587238784\pi^{12}+14680941405810341458359816576\pi^{14}-11947937673613099034164443750\pi^{16}\Big],$$

$$\hat{Z}^{(18,19)}(2)=\frac{1}{12519280390862596113350760561314365440000000\pi^{18}}\Big[-28378559125051743928320000+1565466573304371781435392000\pi^2-362010479258874478428684288000\pi^4$$
$$+37474342135788621074769838080\pi^6-20999466816958669749870646886400\pi^8+66884541720204014704157441126400\pi^{10}-11982818897222541532284369726464\pi^{12}$$
$$+1122064395505490354246756844710\pi^{14}-44890987186261886713204771350000\pi^{16}+35143105400316434035632304687500\pi^{18}\Big],$$

$$\hat{Z}^{(19,20)}(2)=-\frac{1}{18077840884405588787678498250537943695360000000\pi^{19}}\Big[260034050935690604052480000+167378576740920004904091648000\pi^2-46032139411467789197102284800000\pi^4$$
$$+49864936569429230001889571635200\pi^6-29262627461355462454511634923520000\pi^8+10223370259001222313696112466841600\pi^{10}-21126499458367808558188789817036800\pi^{12}$$
$$+2341691134873926453650025102600576\pi^{14}-10758300301892926120908011549919840\pi^{16}+87057436298005995587368943405625\pi^{18}\Big],$$

$$\hat{Z}^{(20,21)}(2)=\frac{1}{2892454541504894206028559720086070991257600000000\pi^{20}}\Big[25690000707001171197296640000-2488551723420168247474913280000000\pi^2+75566817095464546521607294156800\pi^4$$
$$-1062160217433242638061350551552000\pi^6+83475984203142035463930152647065600\pi^8-38893078089971602450053653713305600\pi^{10}+10875548721335722097675694674638131200\pi^{12}$$
$$-1775932910388692220449035532375705600\pi^{14}+15584388998302767740765296288584075840\pi^{16}-59778729021517054930386140592303000\pi^{18}+46306312830726949307050906271484375\pi^{20}\Big].$$

$$\hat{Z}^{(1,2)}{}_{(3)}=\tfrac{1}{12}\big(2\sqrt{3}-3\big),\qquad \hat{Z}^{(2,3)}{}_{(3)}=\tfrac{1}{432}\Big(-27+14\sqrt{3}+\tfrac{9}{\pi}\Big),\qquad \hat{Z}^{(3,4)}{}_{(3)}=-\tfrac{45+18\sqrt{3}-14\sqrt{3}\pi}{1728\pi},\qquad \hat{Z}^{(4,5)}{}_{(3)}=\frac{702+84\big(27+2\sqrt{3}\big)\pi+\big(1152\sqrt{3}-2881\big)\pi^2}{248832\pi^2},$$

$$\hat{Z}^{(5,6)}{}_{(3)}=\frac{54\big(14\sqrt{3}-37\big)+840\sqrt{3}\pi+\big(5797-3574\sqrt{3}\big)\pi^2}{995328\pi^2},\qquad \hat{Z}^{(6,7)}{}_{(3)}=\frac{17982+162\big(182\sqrt{3}-1647\big)\pi+27\big(2304\sqrt{3}-5905\big)\pi^2-7\big(42110\sqrt{3}-78327\big)\pi^3}{322486272\pi^3},$$

$$\hat{Z}^{(7,8)}{}_{(3)}=\frac{-2430\big(61+18\sqrt{3}\big)+83916\sqrt{3}\pi+27\big(28553+16770\sqrt{3}\big)\pi^2+\big(78732-335594\sqrt{3}\big)\pi^3}{1289945088\pi^3},$$

$$\hat{Z}^{(8,9)}{}_{(3)}=\frac{1472580+9072\big(2295+74\sqrt{3}\big)\pi+324\big(91008\sqrt{3}-105709\big)\pi^2-168\big(793233+60254\sqrt{3}\big)\pi^3+\big(178071703-76499136\sqrt{3}\big)\pi^4}{371504185344\pi^4},$$

$$\hat{Z}^{(9,10)}{}_{(3)}=\frac{2916\big(774\sqrt{3}-2857\big)+5533920\sqrt{3}\pi-324\big(152814\sqrt{3}-521209\big)\pi^2-48\big(1483097\sqrt{3}-511758\big)\pi^3+\big(226863738\sqrt{3}-370195279\big)\pi^4}{1486016741376\pi^4},$$

$$\hat{Z}^{(10,11)}{}_{(3)}=\frac{299312820+72900\big(7070\sqrt{3}-171747\big)\pi+24300\big(382464\sqrt{3}-411157\big)\pi^2-2700\big(6698762\sqrt{3}-62488989\big)\pi^3-9\big(5065099200\sqrt{3}-9212744479\big)\pi^4+25\big(6475592722\sqrt{3}-11826421389\big)\pi^5}{4012245201715200\pi^5},$$

$$\hat{Z}^{(11,12)}{}_{(3)}=\frac{-131220\big(28925+6642\sqrt{3}\big)+2915854200\sqrt{3}\pi+24300\big(2404421+1236762\sqrt{3}\big)\pi^2-5400\big(8730505\sqrt{3}-7125246\big)\pi^3-9\big(40242657395+27610808958\sqrt{3}\big)\pi^4+50\big(3671699105\sqrt{3}-1298211948\big)\pi^5}{16048980806860800\pi^5},$$

$$\hat{Z}^{(12,13)}{}_{(3)}=\frac{1}{6933159708563865600\pi^6}\Big[25651672920+3674160\big(281907+4562\sqrt{3}\big)\pi+218700\big(11516544\sqrt{3}-6754681\big)\pi^2-226800\big(106545861+3385430\sqrt{3}\big)\pi^3-162\big(232246756800\sqrt{3}-175868541043\big)\pi^4$$
$$+36\big(3302763448131+214002197506\sqrt{3}\big)\pi^5+25\big(2873091390912\sqrt{3}-6479908382207\big)\pi^6\Big],$$

$$\hat{Z}^{(13,14)}{}_{(3)}=\frac{1}{2773263883425546240\pi^6}\Big[787320\big(66366\sqrt{3}-316045\big)+212550156000\sqrt{3}\pi-218700\big(12640806\sqrt{3}-69160621\big)\pi^2-64800\big(147495509\sqrt{3}-104910390\big)\pi^3$$
$$+162\big(296550133938\sqrt{3}-1076243018035\big)\pi^4+360\big(205884495833\sqrt{3}-99859338540\big)\pi^5-175\big(1189802574054\sqrt{3}-1946635606421\big)\pi^6\Big],$$

$$\hat{Z}^{(14,15)}{}_{(3)}=\frac{1}{1467611247108799070208007^7}\Big[9762153103080+38578680\big(456134\sqrt{3}-24118263\big)\pi+19289340\big(70712064\sqrt{3}-37224121\big)\pi^2-3572100\big(371449442\sqrt{3}-9412651737\big)\pi^3$$
$$-23814\big(1164740241600\sqrt{3}-760466097211\big)\pi^4+7938\big(4920941754202\sqrt{3}-40985286126129\big)\pi^5+9\big(10577414304217152\sqrt{3}-16694829965655623\big)\pi^6-1225\big(241451318806186\sqrt{3}-436239059157621\big)\pi^7\Big],$$

$$\hat{Z}^{(15,16)}{}_{(3)}=\frac{1}{5870444988435196280832007^7}\Big[-49601160\big(3662825+677322\sqrt{3}\big)+170696384888400\sqrt{3}\pi+19289340\big(210402593+126703602\sqrt{3}\big)\pi^2-7144200\big(729601789\sqrt{3}-1620343926\big)\pi^3$$
$$-23814\big(2803076569895+2326367522214\sqrt{3}\big)\pi^4+79380\big(992950495129\sqrt{3}-1601735224140\big)\pi^5+9\big(65702735219040679+49510164883503726\sqrt{3}\big)\pi^6-2450\big(134862880599065\sqrt{3}-58222941612138\big)\pi^7\Big],$$

$$\hat{Z}^{(16,17)}(3)=\frac{1}{33813763133386730577592320 0\pi^8}\Big[873064837174320+1851776640\big(44415567+393626\sqrt{3}\big)\pi+154314720\big(1916774784\sqrt{3}-606477925\big)\pi^2-240045120\big(17798150169+279968270\sqrt{3}\big)\pi^3$$
$$-285768\big(39244249540800\sqrt{3}-13493732619847\big)\pi^4+127008\big(536988724735743+6895335985354\sqrt{3}\big)\pi^5+216\big(518491284724453824\sqrt{3}-326463576135631595\big)\pi^6$$
$$-336\big(893382789200775309+46813383823326550\sqrt{3}\big)\pi^7-1225\big(152389841509146240\sqrt{3}-337119938695893131\big)\pi^8\Big],$$

$$\hat{Z}^{(17,18)}(3)=\frac{1}{13525505253354692231036928 00\pi^8}\Big[1488034800\big(1521342\sqrt{3}-8972381\big)+13565467542816000\sqrt{3}\pi-154314720\big(1439560350\sqrt{3}-11885783617\big)\pi^2$$
$$-68584320\big(20996707697\sqrt{3}-26100169758\big)\pi^3+1428840\big(6210164253042\sqrt{3}-37135089788099\big)\pi^4+1270080\big(26266226178293\sqrt{3}-26656967579580\big)\pi^5$$
$$-216\big(598709018914049250\sqrt{3}-2213891956469549207\big)\pi^6-672\big(304318107429747331\sqrt{3}-172874144162115774\big)\pi^7+1225\big(435235968952169778\sqrt{3}-713337768334931003\big)\pi^8\Big],$$

$$\hat{Z}^{(18,19)}(3)=\frac{1}{13146791106260760848567894016 00\pi^9}\Big[58464585220935600+18749238480\big(5867234\sqrt{3}-600328125\big)\pi+1785641760\big(15138920448\sqrt{3}-4236518269\big)\pi^2$$
$$-462944160\big(31133608058\sqrt{3}-1722444578037\big)\pi^3-115736040\big(10989313693632\sqrt{3}-3280921971535\big)\pi^4+2571912\big(455517813319858\sqrt{3}-7391659237811325\big)\pi^5$$
$$+5832\big(2873264941254076992\sqrt{3}-1573633144207062371\big)\pi^6-13608\big(1569723702002837758\sqrt{3}-1168669161611556644 7\big)\pi^7$$
$$-81\big(6050229328824219843 84\sqrt{3}-886168883247494751971\big)\pi^8+1225\big(114549503631688611466\sqrt{3}-20547051240856276664 1\big)\pi^9\Big],$$

$$\hat{Z}^{(19,20)}(3)=\frac{1}{52587164425043043394271576064 00\pi^9}\Big[-2678462640\big(575940725+87843042\sqrt{3}\big)+1682253111024208800\sqrt{3}\pi+1785641760\big(9018837149+16522525962\sqrt{3}\big)\pi^2$$
$$-925888320\big(64084310353\sqrt{3}-436522842414\big)\pi^3-949035528\big(1252129132998\sqrt{3}-302840637145\big)\pi^4+25719120\big(63997148204653\sqrt{3}-458742838215420\big)\pi^5$$
$$+5832\big(28141886829345656 11+428444048664306523 8\sqrt{3}\big)\pi^6-27216\big(1228646175500522483\sqrt{3}-2983446322924960494\big)\pi^7-81\big(31741120502649059011 51+25568508251916629074 62\sqrt{3}\big)\pi^8$$
$$+2450\big(63005382101415209525\sqrt{3}-30828591348338574072\big)\pi^9\Big],$$

$$\hat{Z}^{(20,21)}(3)=\frac{1}{18931379193015495621937767383040000\pi^{10}}\Big[272125082086312392 00+624974616000\big(8466191307+43655330\sqrt{3}\big)\pi+66961566000\big(390117511296\sqrt{3}-69963418417\big)\pi^2$$
$$-138883248000\big(3538375943949+30304795526\sqrt{3}\big)\pi^3-231472080\big(8146846411531200\sqrt{3}-1434108747628219\big)\pi^4-771573600\big(283388114263510\sqrt{3}-21213224962217631\big)\pi^5$$
$$+437400\big(1032106912553345166 72\sqrt{3}-29179474844566192319\big)\pi^6+1360800\big(1852787727580925522\sqrt{3}-161730054824069504817\big)\pi^7$$
$$-486\big(7700134593283814799648 00\sqrt{3}-4397712665446743999654 23\big)\pi^8+2100\big(4325708010109432379654 97+1759686651160009422895 0\sqrt{3}\big)\pi^9$$
$$+214375\big(2686006964880752857344\sqrt{3}-586172470372204012 0369\big)\pi^{10}\Big].$$

$$\hat{Z}^{(1,2)}(4)=\frac{\pi-2}{16\pi},\quad \hat{Z}^{(2,3)}(4)=\frac{12+12\pi-5\pi^2}{512\pi^2},\quad \hat{Z}^{(3,4)}(4)=\frac{-168+396\pi+202\pi^2-99\pi^3}{73728\pi^3},\quad \hat{Z}^{(4,5)}(4)=\frac{1200+4320\pi-3512\pi^2-4872\pi^3+1755\pi^4}{4718592\pi^4},$$

$$\hat{Z}^{(5,6)}(4)=\frac{-38880+241200\pi+186000\pi^2-400200\pi^3-203494\pi^4+96975\pi^5}{1887436800\pi^5},\quad \hat{Z}^{(6,7)}(4)=\frac{953280+8320320\pi-7378800\pi^2-36784800\pi^3+17373764\pi^4+27667476\pi^5-9333225\pi^6}{543581798400\pi^6},$$

$$\hat{Z}^{(7,8)}(4)=\frac{-52536960+691346880\pi+566479200\pi^2-2914304400\pi^3-2014346488\pi^4+3962357364\pi^5+2156964930\pi^6-995722875\pi^7}{426168129945600\pi^7},$$

$$\hat{Z}^{(8,9)}(4)=\frac{478759680+8468167680\pi-7157041920\pi^2-89293397760\pi^3+38961966624\pi^4+232256453184\pi^5-82822457776\pi^6-145218219408\pi^7+47021834475\pi^8}{54549520633036800\pi^8},$$

$$\hat{Z}^{(9,10)}(4)=\frac{1}{235653929134718976 00\pi^9}\Big[-12959654400+320811321600\pi+249167439360\pi^2-2406136078080\pi^3-1813794333120\pi^4+7622732486880\pi^5$$

$$+5866548067808\pi^6-10329554789424\pi^7-6075970569810\pi^8+2721498152625\pi^9\Big],$$

$$\hat{Z}^{(10,11)}(4)=\frac{1}{1885231433077751808000 0\pi^{10}}\Big[646656998400+20855511936000\pi-15908447520000\pi^2-423480742272000\pi^3+155455887162240\pi^4+2407085602588800\pi^5$$

$$-690712514324000\pi^6-4858102787889600\pi^7+1434686348402316\pi^8+272031066405630 0\pi^9-855380089265625\pi^{10}\Big],$$

$$\hat{Z}^{(11,12)}(4)=\frac{1}{3649808054438527500288000 0\pi^{11}}\Big[-71248933324800+3066836963097600\pi+2147272497216000\pi^2-32994976801248000\pi^3-26307684678401280\pi^4+169824342336485760\pi^5$$

$$+173665340769940800\pi^6-543644538181826400\pi^7-506552450721933352\pi^8+791312771801094444\pi^9+502106969790796050\pi^{10}-218816278991454375\pi^{11}\Big],$$

$$\hat{Z}^{(12,13)}(4)=\frac{1}{2102289439356591840165888000 00\pi^{12}}\Big[2305385523302400+126291787634073600\pi-84666760738560000\pi^2-4394402461709568000\pi^3+1299328657279107840\pi^4$$

$$+4462284259031993856 0\pi^5-9109124891322297600\pi^6-18733351216357261440 0\pi^7+38674141099980946736\pi^8+32414799629592335894 4\pi^9-82963880513737784280\pi^{10}$$

$$-16838145018836223300 0\pi^{11}+51759053721397378125\pi^{12}\Big],$$

$$\hat{Z}^{(13,14)}(4)=\frac{1}{5684590644020224335808561152000 00\pi^{13}}\Big[-326696029973913600+2311074677740308480 0\pi+1422221755947632640 0\pi^2-2965941324152142336 00\pi^3-2627357400324642585 60\pi^4$$

$$+1724522555482742695680\pi^5+3140617642113715146240\pi^6-9346414694706594236160\pi^7-1767587028975945443094 4\pi^8+3840269234571916127467 2\pi^9-4549375668567767917089 6\pi^{10}$$

$$-6304327269971616176522 4\pi^{11}-4297687104962919234465 0\pi^{12}+1827236979240428318062 5\pi^{13}\Big],$$

$$\hat{Z}^{(14,15)}(4)=\frac{1}{8913438129823711758547823886336000 00\pi^{14}}\Big[2639191301216870400 0+2332482753805940736000\pi-135503272345344583680 0\pi^{2}-129495692619143333683200\pi^{3}$$
$$+3004614420021675901440 0\pi^{4}+2094928031999642929689600\pi^{5}-25349601252873628418688 0\pi^{6}-15100547205665994294942720\pi^{7}+16469139979903469748489 60\pi^{8}$$
$$+5265130134094989397503264 0\pi^{9}-80509489974424450171479 52\pi^{10}-81998763263694004366967328\pi^{11}+18677773839866223207584700\pi^{12}+40305044547604234787287500\pi^{13}$$
$$-12163520270564951460046875\pi^{14}\Big],$$

$$\hat{Z}^{(15,16)}(4)=\frac{1}{3208837726736536233077216599080960000 00\pi^{15}}\Big[-46337027028582850560 00+5161133892265660108800 00\pi+2753719229717584128000 00\pi^{2}-62224204317351966489600 00\pi^{3}$$
$$-71456389135528489956864 00\pi^{4}-26932558065808670795520 00\pi^{5}+15454735079658896365344000 0\pi^{6}+15965277339324365522870400 0\pi^{7}-14734877438663429088788947 20\pi^{8}$$
$$+9359041779134835254458992 00\pi^{9}+66858517669427198201851000 00\pi^{10}-96224314067529306641233524 00\pi^{11}-15115257648023669929267894152\pi^{12}+18658828832122713811445030700\pi^{13}$$
$$+13606289597410394667282123750\pi^{14}-56576325163404492016977656 25\pi^{15}\Big],$$

$$\hat{Z}^{(16,17)}(4)=\frac{1}{8214624580445532756677674493647257600000 0\pi^{16}}\Big[56772848827620237312000+7781492578357428682752000\pi-38870522009448114585600 00\pi^{2}-65778295774187062370304000 0\pi^{3}$$
$$+11761902703752893052641280 0\pi^{4}+15734914034909020252626124800\pi^{5}-57401417029713205626777600 0\pi^{6}-17464142127393351033522278400 0\pi^{7}+85299951909826774209433344 0\pi^{8}$$
$$+10195825579040269533266379878 40\pi^{9}-41099081861260979377311526400\pi^{10}-31294693280855809903105478016 00\pi^{11}+35385493289581432311492780230 4\pi^{12}$$
$$+45101798351849695122898694634 24\pi^{13}-93012800845656347128668672720 0\pi^{14}-21238859242627612757121648600 00\pi^{15}+63113992880089590111637026562 5\pi^{16}\Big],$$

$$\hat{Z}^{(17,18)}(4)=\frac{1}{3798442405998014346687756685862491914240000000\pi^{17}}\Big[-12013213941558716350464000+2036573414071306950647808000\pi+93502249631936379666432000 0\pi^{2}$$
$$-12425536068551131275509760000\pi^{3}-32428640849237305307924889600\pi^{4}-78511583802716229255722311680 0\pi^{5}+13574341013424725578593669120 00\pi^{6}+13228798815940219457808006144000\pi^{7}$$
$$-20719602189385098881987341716480\pi^{8}-73298297241245350647746631498240\pi^{9}+14476836275037732758483445895680 0\pi^{10}+82890136643185050776939034201600\pi^{11}$$
$$-53316192474032898335403138355340 8\pi^{12}+48655571129556447419371821745785 6\pi^{13}+10668956513327066384066045289456 00\pi^{14}-11801930009741885894011496400600 00\pi^{15}$$
$$-91686744719217598396068469153125 0\pi^{16}+37363890154955657099344076601562 5\pi^{17}\Big],$$

$$\hat{Z}^{(18,19)}(4)=\frac{1}{32818542387822843955382217765851930139033600000\pi^{18}}\Big[4658557631177262640988160000+95961974798933629893771264000\pi-4104599430827830637909606400\pi^{2}$$
$$-1195885622508123499647683788800\pi^{3}+16257086680570721683137183744000\pi^{4}+401149261361386900409804454297600\pi^{5}+220123491419162984104185237504000\pi^{6}$$
$$-63865778653360774413933217800192000\pi^{7}-73045479980661020865703244141568000\pi^{8}+562347185180079442719083296281292800\pi^{9}+49584754827108241322443815435955200\pi^{10}$$
$$-28400665433282724852235315553688576000\pi^{11}-42016461450097536061971441723143680\pi^{12}+79355528042724429668602542226484812800\pi^{13}-64556095975146628958789565015676608000\pi^{14}$$
$$-10770341005756088272277605044795029760000\pi^{15}+20351346780539808902090966400844650000\pi^{16}+48998090115695487925429889194442250000\pi^{17}-14368438382832295899554277096173437500\pi^{18}\Big],$$

$$\hat{Z}^{(19,20)}(4)=\frac{1}{1895599008320647466862876898155607484830580736000000\pi^{19}}\Big[-1163069790307875418403635200000+29192296360318816884296417280000\pi+1148618412976383132775184793600\pi^{2}$$
$$+2393205764442822539807891374080000\pi^{3}-5131691274212143600707762094080000\pi^{4}-427438453816924891505605959598080000\pi^{5}+4525124757077057389058534210519040000\pi^{6}$$
$$+9344290338431950122740064535621632000\pi^{7}-10522454557357118268739222148957184000\pi^{8}-9028950896774505741213994757643110400\pi^{9}+10746023163461949104861069204115563520\pi^{10}$$
$$+4156049359657685962440745157323627776000\pi^{11}-57881393600330601611761255722545559040\pi^{12}-66186603676499391299245009164286083840\pi^{13}$$
$$+17854525949559119318114414147348844019200\pi^{14}-96420843472907346704815702438798945344\pi^{15}-32019502450115996910069934769477991756000\pi^{16}$$
$$+31943655953441176617242010859682637450000\pi^{17}+26342822672931578023717259405981146875000\pi^{18}-10540094004422001429157980608352539062500\pi^{19}\Big],$$

$$\hat{Z}^{(20,21)}(4)=\frac{1}{3032958413313035946980603037048971975728929177600000000\pi^{20}}\Big[7890027009727842671990538240000+2380484007269648549678894284800000\pi$$
$$-870094846069703809457430528000000\pi^{2}-427316635043437533171964836249600000\pi^{3}+436968570066904841142048930938880000\pi^{4}+193090124628012493226244016993075200000\pi^{5}$$
$$+309344257389397934776570264166400000\pi^{6}-419353566853385599959711300749475840000\pi^{7}-108995682246502516003641497033589350400\pi^{8}+52045203886873421101066154986054594560000\pi^{9}$$
$$+12822605832813196611503900139321369600000\pi^{10}-39083199328307624131617979444901171712000\pi^{11}-65876073600433899703357604453496282700800\pi^{12}$$
$$+17738495826560783790148968492954437084160000\pi^{13}+10909362164629355169418704132227582272000\pi^{14}-46089961828843644049809596010085475827200\pi^{15}$$
$$+25328535748683448729484308375998661305744\pi^{16}+59611453951362462478969982172558895631760000\pi^{17}-10411097664297284727658253596583168746500\pi^{18}$$
$$-26355433564216579998210255788206467937500\pi^{19}+76392973813974105821930412592541417578125\pi^{20}\Big].$$

$$\hat{Z}^{(1,3)}(4)=-\frac{\pi-4}{16\pi},\quad \hat{Z}^{(2,4)}(4)=-\frac{24+8\pi-5\pi^2}{512\pi^2},\quad \hat{Z}^{(3,5)}(4)=\frac{-480+216\pi+404\pi^2-135\pi^3}{73728\pi^3},\quad \hat{Z}^{(4,6)}(4)=\frac{3648+1920\pi-7280\pi^2-3056\pi^3+1611\pi^4}{4718592\pi^4},$$

$$\hat{Z}^{(5,7)}(4)=\frac{149760-91200\pi-542400\pi^2+290000\pi^3+421388\pi^4-145575\pi^5}{1887436800\pi^5},\quad \hat{Z}^{(6,8)}(4)=-\frac{3985920+2695680\pi-23150400\pi^2-14083200\pi^3+33643016\pi^4+17182584\pi^5-8199225\pi^6}{543581798400\pi^6},$$

$$\hat{Z}^{(7,9)}(4)=\frac{-262563840+195310080\pi+2234400000\pi^2-1536561600\pi^3-6423200672\pi^4+3327482984\pi^5+4550305860\pi^6-1570037175\pi^7}{426168129945600\pi^7},$$

$$\hat{Z}^{(8,10)}(4)=\frac{2612736000+2100510720\pi-30784481280\pi^2-23158732800\pi^3+108930057600\pi^4+86101125376\pi^5-150569774944\pi^6-90240714720\pi^7+40191979275\pi^8}{54549520633036800\pi^8},$$

$$\hat{Z}^{(9,11)}(4)=\frac{1}{2356539291347189760 0\pi^9}\Big[81765089280-70543872000\pi-1275440947200\pi^2+1042115880960\pi^3+836623868774 4\pi^4-496546666560 0\pi^5-19810036547200\pi^6+9535303927008\pi^7$$
$$+12943323415332\pi^8-4424232492225\pi^9\Big],$$

$$\hat{Z}^{(10,12)}(4)=-\frac{1}{1885231433077751808000 0\pi^{10}}\Big[4457056665600+4088254464000\pi-891060065280 00\pi^2-779504947200 00\pi^3+53806372010496 0\pi^4+60459101176320 0\pi^5-17017974146144 00\pi^6$$
$$-18077516530880 00\pi^7+24735167192985 84\pi^8+16965713522754 00\pi^9-71738200764562 5\pi^{10}\Big],$$

$$\hat{Z}^{(11,13)}(4)=\frac{1}{3649808054438527500288000 0\pi^{11}}\Big[-555876886118400+539303856537600\pi+13867726473216000\pi^2-12915778917888000\pi^3-17348466523981824 0\pi^4+10198957099516416 0\pi^5$$
$$+87423013050996480 0\pi^6-41658074727410240 0\pi^7-17594370749387284 16\pi^8+78175312925467586 4\pi^9+10750736136218769 00\pi^{10}-36329808726937687 5\pi^{11}\Big],$$

$$\hat{Z}^{(12,14)}(4)=\frac{1}{2102289439356591840165888000 0\pi^{12}}\Big[19613815190323200+20011567900262400\pi-597928491933696000\pi^2-588280335261696000\pi^3+4783833481684561920\pi^4$$
$$+8202607253775728640\pi^5-234738599925130752 00\pi^6-485814569965705728 00\pi^7+838914533248310794 88\pi^8+122557375894589640576\pi^9-136781388348577452720\pi^{10}$$
$$-105487589166075975600\pi^{11}+428026784218356881 25\pi^{12}\Big],$$

$$\hat{Z}^{(13,15)}(4)=\frac{1}{5684590644020224335808561152000 00\pi^{13}}\Big[3101121451524096000-3314734767164620800\pi-11349201055440568320 0\pi^2+11745553845266841600 0\pi^3+25105699042707419136 00\pi^4$$
$$-12558943050232032344 80\pi^5-22531100208573717995520\pi^6+81453235388666718720 00\pi^7+90366000850395251421440\pi^8-33958009378858482961472\pi^9-15958079308027058500 3968\pi^{10}$$
$$+65614854027384539289360\pi^{11}+92249506159524359154900\pi^{12}-30813026723470622255625\pi^{13}\Big],$$

$$\hat{Z}^{(14,16)}(4)=-\frac{1}{8913438129823711758547823886336000 00\pi^{14}}\Big[272456547603893452800+303909902249361408000\pi-1179632995900873113600 0\pi^2-1277937497215248629760 0\pi^3$$
$$+8840264957586139004928 0\pi^4+3009451549940616388608 00\pi^5-2746525913320048763904 00\pi^6-3040999319565357886095360\pi^7+2392964858229889270559232\pi^8+1437145134611630403511552 0\pi^9$$
$$-1522619942044631547933088 0\pi^{10}-3162413190895552792147622 4\pi^{11}+2963604783093558115603788 0\pi^{12}+2536779939883201265801940 0\pi^{13}-9949768870178118843736875\pi^{14}\Big],$$

$$\hat{Z}^{(15,17)}(4)=\frac{1}{3208837726736536233077216599080960000 00\pi^{15}}\Big[-528165811261569761280 00+6130272321087602688000 0\pi+2671384160433137909760000\pi^2-3019623748856863948800000\pi^3$$
$$-1018011041406925029015552 00\pi^4+3435939404425857750220800 0\pi^5+1479560245436840096065536000\pi^6-2466078116347960187673600 00\pi^7-9938438552495431469565634560\pi^8$$
$$+1932188409365037474069619200\pi^9+3332097070865404929060122240 0\pi^{10}-9815668497514759021394616000\pi^{11}-5297734414635755446516958985 6\pi^{12}+2025259234292601553953745980 0\pi^{13}$$
$$+2923469796629746447370484750 0\pi^{14}-9655969846089559832542828125\pi^{15}\Big],$$

$$\hat{Z}^{(16,18)}(4)=\frac{1}{8214624580445532756677674493647257600000 00\pi^{16}}\Big[7016881101916361195520 00+8450652980185116180480 00\pi-4101529230079423414272000 0\pi^2-4821252480741871190016000 0\pi^3$$
$$+3337755474698738694881280 0\pi^4+1883902643769674227462963200\pi^5+7405301480379404495781888000\pi^6-3001130749308280732080537600 0\pi^7-5803663628913688335597735936 0\pi^8$$
$$+2288636139339174541242313640960\pi^9+6459190489552270356353259520 0\pi^{10}-9020960397317720988426128384 00\pi^{11}+5661882748170492265142361469 44\pi^{12}+1775528104008903044065560976896\pi^{13}$$
$$-1427367442670304033284940916800\pi^{14}-1342811627750294054504431560000\pi^{15}+5117887037406938492304012656 25\pi^{16}\Big],$$

$$\hat{Z}^{(17,19)}(4)=\frac{1}{3798442405998014346687756685862491914240000000\pi^{17}}\Big[16264367202354883618406400 0-2027878638453828385505280 00\pi-1088345015336053027897344000 0\pi^2$$
$$+1327073843556498740871168000 0\pi^3+7132900977417068845194018816 00\pi^4-8655713620196816043096145920 0\pi^5-1574494265244780316697650790400 0\pi^6-1036043702904303284818378752000\pi^7$$
$$+1613224081182128694869911678156 80\pi^8+6514261302033641531650754519040\pi^9-8722043441151419925811586420736 00\pi^{10}+6469712714389533219441793464320 0\pi^{11}$$
$$+2544365487742659413806808205145088\pi^{12}-5834881514206078398420584604572 16\pi^{13}-3713942049814044603212945067705600\pi^{14}+1326599155381451599614802920984000\pi^{15}$$
$$+1970134581088094635028490942262500\pi^{16}-6438544351172098563033314585156 25\pi^{17}\Big],$$

$$\hat{Z}^{(18,20)}(4)=\frac{1}{32818542387822843955382217765851930139033600000\pi^{18}}\Big[-6818632796571774159421440000-878275828927163715393945600\pi+5180450915471143295558615040\pi^{2}$$
$$+65364675348585309662085120000\pi^{3}+976706133287292305413138022400\pi^{4}-426382456246235210340907352064\pi^{5}-514541701082903817796128630374400\pi^{6}$$
$$+10155450490142756998457964625920\pi^{7}+650537221735844501137655312056320\pi^{8}-1153314059483324470082229627420672\pi^{9}-342697647534207978288414739868467\pi^{10}$$
$$+706740199502358843459426383142912\pi^{11}+605992703720936040856662625211699\pi^{12}-241041349850696485773413550483658752\pi^{13}$$
$$+80659913268744637454130425647020288\pi^{14}+43255430254488446840416011095432448\pi^{15}-30322174834748589115933525656633300\pi^{16}$$
$$-31111760763281724696338752285537500\pi^{17}+11568116317100784064519906346273437\pi^{18}\Big],$$

$$\hat{Z}^{(19,21)}(4)=-\frac{1}{1895599008320647466862876898155607484830580736000000\pi^{19}}\Big[1853195352508743683514826752000-24615264395624104715511398400\pi-15868702255552000608750796800000\pi^{2}$$
$$+20678001113751773824318596710400\pi^{3}+1801900062430180718029237943009280\pi^{4}+21163675594143702518225531043840\pi^{5}-57344396851812489192163049000140800\pi^{6}$$
$$-17045047562763786782105673140797440\pi^{7}+842727817537850131760449924209967104\pi^{8}+233209410540540137287921279309086720\pi^{9}-67491543737021248955405868724821196800\pi^{10}$$
$$-11719930691169437550192833629694238720\pi^{11}+3103746388578388715903304382465607843840\pi^{12}+455124148160941442287835006370644992\pi^{13}-81133647399064932435681871122849618432\pi^{14}$$
$$+14295447854225934774185883539923763788\pi^{15}+1103438222962196441806408655034632274336\pi^{16}-369962673389562437047259920111271157000\pi^{17}$$
$$-56576181841823777772341557535666469750\pi^{18}+18306469686857626753225286629234457812\pi^{19}\Big],$$

$$\hat{Z}^{(20,22)}(4)=\frac{1}{30329584133130359469806030370489719757289291776000000000\pi^{20}}\Big[135524964632671292532342128640000+185319535250874368351482675200000\pi$$
$$-12994366429554983728012748390400000\pi^{2}-1745399012676553111503686860800000\pi^{3}-826703439458812008006079387533312000\pi^{4}$$
$$+19298811917362659740304655830220800000\pi^{5}+44706162801630300252375901997629440000\pi^{6}-65718247693094483918733454344192000000\pi^{7}$$
$$-79906175511141139198225382283328880640\pi^{8}+10540343654758637026632833622300426240000\pi^{9}+69969753605564205304210619610322206720\pi^{10}$$
$$-93397686408797764702005242640750182400\pi^{11}-31113410956395586861188424176997444116480\pi^{12}+484059143019878626640707879177460178944\pi^{13}$$
$$+57217667515278489114848713777537890611200\pi^{14}-14693805000190193465503764702414236620800\pi^{15}+18745796253838016877363977524061318956224\pi^{16}$$
$$+24398410876176559279993564299341952121920\pi^{17}-15106824734124773270840677236727292997000\pi^{18}$$
$$-16802179738768659620772836355703700925000\pi^{19}+61136238207447676268730807457781098828125\pi^{20}\Big].$$

$$\hat{Z}^{(1,2)}(6)=\frac{9-\sqrt{3}\pi}{108\pi},\quad \hat{Z}^{(2,3)}(6)=-\frac{432+408\sqrt{3}\pi-269\pi^2}{31104\pi^2},\quad \hat{Z}^{(3,4)}(6)=\frac{-9720+12960\sqrt{3}\pi+4347\pi^2-1931\sqrt{3}\pi^3}{10077696\pi^3},\quad \hat{Z}^{(4,5)}(6)=\frac{536544+1441152\sqrt{3}\pi-557280\pi^2-3523152\sqrt{3}\pi^3+1912867\pi^4}{58047528896\pi^4},$$

$$\hat{Z}^{(5,6)}(6)=\frac{87130080-307346400\sqrt{3}\pi+310165200\pi^2+911260800\sqrt{3}\pi^3+70893099\pi^4-108143075\sqrt{3}\pi^5}{15672832819200\pi^5},$$

$$\hat{Z}^{(6,7)}(6)=-\frac{5391567360+29989543680\sqrt{3}\pi+47373919200\pi^2-245744668800\sqrt{3}\pi^3-93947321232\pi^4+416879149464\sqrt{3}\pi^5-207252738175\pi^6}{13541327555788800\pi^6},$$

$$\hat{Z}^{(7,8)}(6)=-\frac{1525334872320-10774871009280\sqrt{3}\pi+44069352052320\pi^2+87765668272800\sqrt{3}\pi^3-162341159067864\pi^4-195569135070624\sqrt{3}\pi^5+2889114383061\pi^6+21333980075275\sqrt{3}\pi^7}{716607054252343296 00\pi^7},$$

$$\hat{Z}^{(8,9)}(6)=\frac{1}{8255313264986994769920 0\pi^8}\Big[103900541875200+1038163124920320\sqrt{3}\pi+56082956699750 40\pi^2-18539127946644480\sqrt{3}\pi^3-5005433103973344 0\pi^4+93338872774430976\sqrt{3}\pi^5$$

$$+62169932532485952\pi^6-135509099423062944\sqrt{3}\pi^7+63809267778892475\pi^8\Big],$$

$$\hat{Z}^{(9,10)}(6)=\frac{1}{80241644935673589163622400\pi^9}\Big[4888772865930240-6064742245556736 0\sqrt{3}\pi+5563218044060006 40\pi^2+933542468377989120\sqrt{3}\pi^3-5164192759594454208\pi^4$$

$$-5149624432989572928\sqrt{3}\pi^5+1223622341645592643 2\pi^6+1079398967281351833 6\sqrt{3}\pi^7-712918751457709125\pi^8-1128948143041047875\sqrt{3}\pi^9\Big],$$

$$\hat{Z}^{(10,11)}(6)=-\frac{1}{5777398435368498419780812800 00\pi^{10}}\Big[1808196272496230400+29820491666574336000\sqrt{3}\pi+341976908820339072000\pi^2-95667186503792332 8000\sqrt{3}\pi^3-6334631015171771612160\pi^4$$

$$+96782446177267229568 00\sqrt{3}\pi^5+30714971087435892321600\pi^6-39110535608560387142400\sqrt{3}\pi^7-32542174192979041907184\pi^8+52142498074569979739400\sqrt{3}\pi^9-23628686735164823781875\pi^{10}\Big],$$

$$\hat{Z}^{(11,12)}(6)=-\frac{1}{7549904275339553734969566167040000\pi^{11}}\Big[1050190755685895577600-21133567503116999884800\sqrt{3}\pi+353851100633289764736000\pi^2+5120584409475376047360 00\sqrt{3}\pi^3$$

$$-5892361026837026926947840\pi^4-4862552393702426691502080\sqrt{3}\pi^5+3440752924024899624984480 0\pi^6+2436185002018643776406880 0\sqrt{3}\pi^7-7013789493503923362345933 6\pi^8$$

$$-5194686715624312708544899 2\sqrt{3}\pi^9+5204432021907025587870225\pi^{10}+5303975247315120333409375\sqrt{3}\pi^{11}\Big],$$

$$\hat{Z}^{(12,13)}(6)=\frac{1}{1304623458778674885402741033664512000 0\pi^{12}}\Big[8349081573517296353280 0+2145434180984613421056000\sqrt{3}\pi+4366807743686046991073280 0\pi^2-1107458662245876736266240 00\sqrt{3}\pi^3$$

$$-1387778216242127674344145920\pi^4+1882767167057484541106380800\sqrt{3}\pi^5+1380568969188016983190176768 0\pi^6-1459441842490028799217221120 0\sqrt{3}\pi^7-5139894852274243578166273885 28\pi^8$$

$$+5224727166861231046269299520 0\sqrt{3}\pi^9+4965216196413700770540908515 2\pi^{10}-6609538231170706429586600760 0\sqrt{3}\pi^{11}+2908585445449518298225699437 5\pi^{12}\Big],$$

$$\hat{Z}^{(13,14)}(6)=\frac{1}{2381198736962837400837082934644467302400 00\pi^{13}}\Big[6284926699640314280755200 0-1943173953584639751809433600\sqrt{3}\pi+5338595844457204110339317760 0\pi^2$$

$$+6602357815237950899668254720 0\sqrt{3}\pi^3-1363985460563660248877968204800\pi^4-8772948727855909565603156121 60\sqrt{3}\pi^5+1400500268091950234621259234816 0\pi^6$$

$$+7269144146566425060461610577920\sqrt{3}\pi^7-6850751225041228942727603110512 0\pi^8-3816405706008343536628554685024\sqrt{3}\pi^9+1306995224051403708548515212284 64\pi^{10}$$

$$+8614621976844329771624780611852 8\sqrt{3}\pi^{11}-1079654762193537068091666208462 5\pi^{12}-8667048310788410692867517174375\sqrt{3}\pi^{13}\Big],$$

$$\hat{Z}^{(14,15)}(6)=-\frac{1}{33603476576019561400612914373702722571468800000\pi^{14}}\Big[372807001724637795342483456000+1430800961782505255711716147200\sqrt{3}\pi+469595319744883200878216503296000\pi^{2}$$
$$-1107345094119353925068021268480000\sqrt{3}\pi^{3}-2318704455841811380006441016279040\pi^{4}+2843694054617147244933909889720320\sqrt{3}\pi^{5}+3888304124515878659997729675326976 0\pi^{6}$$
$$-3559299624608229907071131724779520 0\sqrt{3}\pi^{7}-2874793613965990573097761042354191360\pi^{8}+2383288241901898880150578117409748 48\sqrt{3}\pi^{9}+9134524268696410200306607436491976 64\pi^{10}$$
$$-7937010249234225074905639138903814 40\sqrt{3}\pi^{11}-8285028049201318602580877045789432 16\pi^{12}+9700101730886740068237379014053706 00\sqrt{3}\pi^{13}-4169237970969802695969226090449831 25\pi^{14}\Big],$$

$$\hat{Z}^{(15,16)}(6)=-\frac{1}{272188160265758447344964606426992052828897280000000\pi^{15}}\Big[116555811361098330311681310720 00-5319457232706339747883175116800 00\sqrt{3}\pi$$
$$+2245934674718333488804473022464000 0\pi^{2}+2325965140144114274335545759744000 0\sqrt{3}\pi^{3}-7986495711104624141223653099354112 00\pi^{4}-3540386687001097497091510390086 80000\sqrt{3}\pi^{5}$$
$$+1228585428928712188396409195270822400 0\pi^{6}+3605043505396224389483991099878784000\sqrt{3}\pi^{7}-1039492524722700338770526841263550681 60\pi^{8}-3434879007911382083169582875360670 7200\sqrt{3}\pi^{9}$$
$$+4711391377094140895293786101574144296 00\pi^{10}+2156182505654760733041429013373480856 00\sqrt{3}\pi^{11}-8737851775637996259002386124488087 49496\pi^{12}$$
$$-5288042137951996758484679352081306024 00\sqrt{3}\pi^{13}+7669918275563632845342730483639577812 5\pi^{14}+5270641579036823042015246885255342187 5\sqrt{3}\pi^{15}\Big],$$

$$\hat{Z}^{(16,17)}(6)=\frac{1}{6271215212523074626827984530277896897177793331200000\pi^{16}}\Big[104578798782418783905625237094400 0+5801412931331279405685937274880000 0\sqrt{3}\pi$$
$$+2890261326910604797067083799592960000\pi^{2}-6413144343783076626904574440243200000\sqrt{3}\pi^{3}-2082615272049804478643555972810440704 00\pi^{4}+2312208443326366646490270398425989120 00\sqrt{3}\pi^{5}$$
$$+5310167958631405213406042497343963136000\pi^{6}-4199879795694261520913887713026088960000\sqrt{3}\pi^{7}-6463603365450938874766346827067699622912 0\pi^{8}$$
$$+4447637993835217569435276885444824555520 0\sqrt{3}\pi^{9}+3975551352993623067661115604587586591744 00\pi^{10}-2742145114097054270657441067082154150400 00\sqrt{3}\pi^{11}$$
$$-1133224637471468481853392175295954525552691 2\pi^{12}+8738839805425357983864716388416396999500 80\sqrt{3}\pi^{13}+9809236552306120557255055048677433287840 00\pi^{14}$$
$$-1043126836337282812319811534016246860360000\sqrt{3}\pi^{15}+4396549991659857127849474089270953519218 75\pi^{16}\Big],$$

$$\hat{Z}^{(17,18)}(6)=\frac{1}{19573716921327020525255505321521531795471328545341440000000\pi^{17}}\Big[11871886165001881047463916678676480 00-7753613365833491150155422877129113600 0\sqrt{3}\pi$$
$$+4810739638569891348192726073877987328000\pi^{2}+4054475696913622311424902227070812160000\sqrt{3}\pi^{3}-2217040367307168527437878895312110256128 00\pi^{4}$$
$$-4865362099561806562586129851712550666240 0\sqrt{3}\pi^{5}+4593816991830289286895674382544697479987200\pi^{6}+4098855588170186601326377492456085913600 0\sqrt{3}\pi^{7}$$
$$-5723535931474716783142801669386105764926464 0\pi^{8}-1398797694358764418873290184223707594782720\sqrt{3}\pi^{9}+4494861575609451783090229532373314903249817 60\pi^{10}$$
$$+8173600490446058003637400629840055415255040 0\sqrt{3}\pi^{11}-1991149843318171818517305815241134645209551744\pi^{12}-7630948174865257799847762496043092791129918 72\sqrt{3}\pi^{13}$$
$$+3665688600395452658283818679073093240377865536\pi^{14}+2076250465297094073190381210322975790916915200\sqrt{3}\pi^{15}-3346390615198831968541006626376223798495831 25\pi^{16}$$
$$-2057053175376214672418861657867367138549218 75\sqrt{3}\pi^{17}\Big],$$

$$\hat{Z}^{(18,19)}(6)=-\frac{1}{5073507426007963720146226979338381041386168358952501248000000\pi^{18}}\Big[1121696674259079097548279470201241600000+87610096826113273055215365468570255360000\sqrt{3}\pi$$
$$+635332157212957254490147188814503346176000\pi^2-13359785071297239188516825120432141107200000\sqrt{3}\pi^3-639754693933582651062125101288574075535360000\pi^4$$
$$+639803201737028779422520934043526910705664000\sqrt{3}\pi^5+232332610409581608843037049881060808511651840\pi^6-155713487849726599955950349392734917099520000\sqrt{3}\pi^7$$
$$-42141943716694674901775928843034754518482124800\pi^8+2341586347929766275216269154831340360153178112\sqrt{3}\pi^9+4189180672365028711089797170711713200073704755200\pi^{10}$$
$$-22759793546626185224050071508124801873515438080\sqrt{3}\pi^{11}-22667378683950361037656293417421779181074133708800\pi^{12}+13453574497405614340871131712437891556518382008320\sqrt{3}\pi^{13}$$
$$+5963113721796320541990915258671906871421660762752\pi^{14}-417780944640803381869631871054458441031456864000\sqrt{3}\pi^{15}-49769379912301923607349436996137208200247293052000\pi^{16}$$
$$+490574710379871205326025475300136334815377152500\sqrt{3}\pi^{17}-203342164443336911403911141932125566937817548437\pi^{18}\Big],$$

$$\hat{Z}^{(19,20)}(6)=-\frac{1}{19780590752519848952106109747044480004156393197884011865702400000\pi^{19}}\Big[150731009346983096078813746873519570944000-13741022465134014916631041692255538642944000\sqrt{3}\pi$$
$$+12123713005789194773030649344142689772503040000\pi^2+799266787251875790995461175224188516827136000\sqrt{3}\pi^3-6828230458987052379164472787426687969764311040\pi^4$$
$$+26757060809545802040965336686219816435974144\sqrt{3}\pi^5+17395986227198536662665342552558133319240482816\pi^6-667076322839036484948869902342131197617265049600\sqrt{3}\pi^7$$
$$-28241976718243698252529841837165906826591884083\pi^8+1280903411827676457759446163659297525817631997952\sqrt{3}\pi^9+3294580610462715989066062732856594505348915848908\pi^{10}$$
$$-85031268046571243105039459934792110968191825684480\sqrt{3}\pi^{11}-260257162819806864508575680031600217291293185151283\pi^{12}-11851978188988616015440253883053913185984130854092\sqrt{3}\pi^{13}$$
$$+11679507931222724417959690597437582046450926254080128\pi^{14}+377045780671513654885483261277134531716307608084979\sqrt{3}\pi^{15}-21623284602056709093448036211277283146884464073758728\pi^{16}$$
$$-11609532246621221063550632589619964332437599404471200\sqrt{3}\pi^{17}+2029168372762486258176852925611507246328865226208125\pi^{18}+1145894148639381402666892230564021688810808531921875\sqrt{3}\pi^{19}\Big],$$

$$\hat{Z}^{(20,21)}(6)=\frac{1}{28484050683628582491032798035744051205985206204952977086611456000000000\pi^{20}}\Big[7464292546483844954754730478872018550784000+80326442697464330888028910890455842947072000\sqrt{3}\pi$$
$$+821914392151038888548315083775217513686630400000\pi^2-16466278100941183397543330005945766178717696000\sqrt{3}\pi^3-1120721228358451055985957765524123467960917688320\pi^4$$
$$+10007892257232637342638551556463219347840565248000\sqrt{3}\pi^5+553757517655688498124541735364351388494716993536000\pi^6-3038395184049792755100522916786179694453964144640000\sqrt{3}\pi^7$$
$$-14047155889029457438635062711255227302410950625034240\pi^8+59237557362219389544947632545655820631827990544384000\sqrt{3}\pi^9+204926299140023769371505625384384559919626013777100800\pi^{10}$$
$$-81918678794705784139803136316277309842531324140339200\sqrt{3}\pi^{11}-17666867538774157240250172783244591468417805177490094080\pi^{12}$$
$$+77682308142066407842257746495154043718698829997379584\sqrt{3}\pi^{13}+868895953206164834067455433982768449375499030891281408\pi^{14}$$
$$-45333491603535871102721096808975506192752031259363379200\sqrt{3}\pi^{15}-21478128595487145758291415245148376740123587189957069376\pi^{16}$$
$$+13890015055226906890128064629310632794488048163993588160\sqrt{3}\pi^{17}+17405518431688624898189025662140601907512499664111506000\pi^{18}$$
$$-16124114980335880989039065922615314033255691692269125000\sqrt{3}\pi^{19}+65870864016187141559943692468582292090670138827658984375\pi^{20}\Big],$$

$$\hat{Z}^{(1,3)}(6)=\frac{7}{72}-\frac{1}{2\sqrt{3}\pi},\quad \hat{Z}^{(2,4)}(6)=\frac{54+108\sqrt{3}\pi-65\pi^2}{5184\pi^2},\quad \hat{Z}^{(3,5)}(6)=\frac{-4212\sqrt{3}+3969\pi+8010\sqrt{3}\pi^2-4583\pi^3}{1679616\pi^3},\quad \hat{Z}^{(4,6)}(6)=\frac{1944+21600\sqrt{3}\pi-360\pi^2-33024\sqrt{3}\pi^3+17017\pi^4}{35831808\pi^4},$$

$$\hat{Z}^{(5,7)}(6)=\frac{-94828320\sqrt{3}+117369000\pi+661802400\sqrt{3}\pi^2-344725200\pi^3-1082915916\sqrt{3}\pi^4+594333475\pi^5}{5224277606400\pi^5},$$

$$\hat{Z}^{(6,8)}(6)=\frac{7873200+304780320\sqrt{3}\pi+154693800\pi^2-1695427200\sqrt{3}\pi^3-396367290\pi^4+1897354716\sqrt{3}\pi^5-914518625\pi^6}{41794220851200\pi^6},$$

$$\hat{Z}^{(7,9)}(6)=\frac{-508378822560\sqrt{3}+758572584840\pi+7829183038320\sqrt{3}\pi^2+1097913511800\pi^3-39580063351668\sqrt{3}\pi^4+7998307736517\pi^5+61102529757546\sqrt{3}\pi^6-32343075907625\pi^7}{5971725452102860800\pi^7},$$

$$\hat{Z}^{(8,10)}(6)=\frac{1}{7643808578691661824 00\pi^8}\Big[374984769600+40399417336320\sqrt{3}\pi+36931216049280\pi^2-500894881966080\sqrt{3}\pi^3-365736091175280\pi^4+1829353921448256\sqrt{3}\pi^5$$
$$+641781150699024\pi^6-1813331818276224\sqrt{3}\pi^7+839054170880975\pi^8\Big],$$

$$\hat{Z}^{(9,11)}(6)=\frac{1}{133736074892789315272704 00\pi^9}\Big[-3917340893088000\sqrt{3}+6730001653919040\pi+106813099769080320\sqrt{3}\pi^2+195151699634814720\pi^3-109306740361148064 0\sqrt{3}\pi^4$$
$$-1393374237131561328\pi^5+4806535472237733696\sqrt{3}\pi^6+225249339667869216\pi^7-718080089717294089 2\sqrt{3}\pi^8+3687887513094183875\pi^9\Big],$$

$$\hat{Z}^{(10,12)}(6)=\frac{1}{53494429957115726109081600 0\pi^{10}}\Big[546727794076800+142828556011142400\sqrt{3}\pi+176372211457512000\pi^2-317125195817794560 0\sqrt{3}\pi^3-345921903953977680 0\pi^4+240591845834965699 20\sqrt{3}\pi^5$$
$$+21076122335055152400\pi^6-73691721088732385280\sqrt{3}\pi^7-319004238616372294 02\pi^8+68744639166219603180\sqrt{3}\pi^9-308993527972153176 25\pi^{10}\Big],$$

$$\hat{Z}^{(11,13)}(6)=\frac{1}{3145793448058147389570652569600 00\pi^{11}}\Big[-249184525751800531200\sqrt{3}+479113179190351675200\pi+1062656065197837633600 0\sqrt{3}\pi^2+5581484314879459140000 0\pi^3$$
$$-1826211467521600362403 20\sqrt{3}\pi^4-8632829991970510790032 80\pi^5+1549450367229406203856800\sqrt{3}\pi^6+3440470376638837660734000\pi^7-6279315699721705626883788\sqrt{3}\pi^8-1560037420137424782621177\pi^9$$
$$+9188182264304984094662550\sqrt{3}\pi^{10}-4599968240521538521223125\pi^{11}\Big],$$

$$\hat{Z}^{(12,14)}(6)=\frac{1}{2013307806757214329325217644544000 0\pi^{12}}\Big[35723194064978112000+2057476784106607395840 0\sqrt{3}\pi+3082082734674787979520 0\pi^2-7219324364020720680960 00\sqrt{3}\pi^3$$
$$-9922483030855848480720 00\pi^4+9271329452609638568394240\sqrt{3}\pi^5+1187203179093124918076352 0\pi^6-5680383188913900942266880 0\sqrt{3}\pi^7-5757536935884810375930708 0\pi^8$$
$$+1591739214377908335380372 16\sqrt{3}\pi^9+7949083011952728044725048 8\pi^{10}-1435639660939177704710112 00\sqrt{3}\pi^{11}+6310025025495664359092937 5\pi^{12}\Big],$$

$$\hat{Z}^{(13,15)}(6)=\frac{1}{1984332280802364500697569112203722752000 0\pi^{13}}\Big[-3490734779786998854190080 0\sqrt{3}+7369318364492915469999360 0\pi+2149737429734240208774912000\sqrt{3}\pi^2+2483894877525854441242513920 0\pi^3$$
$$-5573821308929028196575101568 0\sqrt{3}\pi^4-6459926148545812602075438422 40\pi^5+7686983972383305878993417856 00\sqrt{3}\pi^6+5582474179276010303493566680320\pi^7-5816270665660415851684740035232\sqrt{3}\pi^8$$
$$-1763788467248502169546483664709 6\pi^9+2236596489290971263617917488720 0\sqrt{3}\pi^{10}+9328711450693810178999289937968\pi^{11}$$
$$-3228188442031365528451232110530 0\sqrt{3}\pi^{12}+1581116125510599137245309836812 5\pi^{13}\Big],$$

$$\hat{Z}^{(14,16)}(6)=\frac{1}{1111226077249324120390638702834084741120000\pi^{14}}\Bigg[293408882133291225100800+34714663661197611511580160000\sqrt{3}\pi+59791006402023881661925248000\pi^{2}$$
$$-17751405274573845408032378880\sqrt{3}\pi^{3}-285984225780228175715064950400\pi^{4}+34152261307985755029067068096000\sqrt{3}\pi^{5}+562268558141996160817557363168\pi^{6}$$
$$-343270730947084149042818701516805\sqrt{3}\pi^{7}-52187593470056431682665384974960\pi^{8}+18912397681602746949953524543344\sqrt{3}\pi^{9}+2181544201447379719272969379452\pi^{10}$$
$$-505688124606911960251521392502912\sqrt{3}\pi^{11}-28168165733688302584721547664578\pi^{12}+4472965259467487408053758202887\sqrt{3}\pi^{13}-193055602699026128461096502430625\pi^{14}\Bigg],$$

$$\hat{Z}^{(15,17)}(6)=\frac{1}{5670586672203300986353429300562334433935360000000\pi^{15}}\Bigg[-1875571281287764093758396672000\sqrt{3}+428836316291516217464059680000\pi$$
$$+157771603497621602127078760320000\sqrt{3}\pi^{2}+3548618715386777350340651428800000\pi^{3}-57548356905014447990024731992768\sqrt{3}\pi^{4}-13899068685170648730622632901140000\pi^{5}$$
$$+11691875593107749555871211732075200\sqrt{3}\pi^{6}+2024984536602142743849987695409480000\pi^{7}-139879198257449176613489888505683664\sqrt{3}\pi^{8}-13394306269662689467865491966274031000\pi^{9}$$
$$+98281871258554167637600853866719468\sqrt{3}\pi^{10}+369139207166201682521912610449237070\pi^{11}-365291251363590051954404252021097931724\sqrt{3}\pi^{12}$$
$$-20531503844503774756475712806217635325\pi^{13}+52250200075877305640284378871036133750\sqrt{3}\pi^{14}-25106273358080443247860845512855921875\pi^{15}\Bigg],$$

$$\hat{Z}^{(16,18)}(6)=\frac{1}{14516701880840450525064779009439576150874521600000\pi^{16}}\Bigg[4990885085087283738964608000+1148069990630487253328755531776000\sqrt{3}\pi$$
$$+22041231774338203776356674867200\pi^{2}-8083216613730421174729525936128000\sqrt{3}\pi^{3}-14687098170983822604244313561472000\pi^{4}+213766270452470808694201332876963840\sqrt{3}\pi^{5}$$
$$+43335078559097330531650025410629120\pi^{6}-3116356148718465202811439940182220800\sqrt{3}\pi^{7}-650276539214003776929407446228214880\pi^{8}+27848184340014788644164145334285787520\sqrt{3}\pi^{9}$$
$$+5070103938533379879253513952880729408\pi^{10}-14615165439046809033424073932154959872\sqrt{3}\pi^{11}-189838370382894027919707296713879698000\pi^{12}$$
$$+38195763629191496977382697770212251467\sqrt{3}\pi^{13}+232547309009543596111752860586810592560\pi^{14}$$
$$-33411774510635689374912609723732427200\sqrt{3}\pi^{15}+14201838907478693115744906156531587656\pi^{16}\Bigg],$$

$$\hat{Z}^{(17,19)}(6)=\frac{1}{81557153838862585521897938839673049147797202272256000000\pi^{17}}\Bigg[-43911298319593670114575897587916800\sqrt{3}+10762785567926988176097668695864320\pi$$
$$+484250657774445180673781031933788160\sqrt{3}\pi^{2}+19720205498260870100087999136279552000\pi^{3}-2365131596008101666098080900747089715200\sqrt{3}\pi^{4}$$
$$-10869474151761264849654859105823141329920\pi^{5}+663738918908137618099762508691944577024\sqrt{3}\pi^{6}+23779984083639268881565933407584164224000\pi^{7}$$
$$-11356209158240507529559937033247458037376\sqrt{3}\pi^{8}-26101912531467412821019263038647372999820160\pi^{9}+12385805906155630070979951640823682035205120\sqrt{3}\pi^{10}$$
$$+14663039556868460687673858217689968877331200\pi^{11}-83256519394360944715825475953694806545127296\sqrt{3}\pi^{12}-36828866133133202605392131750653027670816745 6\pi^{13}$$
$$+30304790015949379342750542743946047418184435 2\sqrt{3}\pi^{14}+20892391590669433404372104332138011460908000 0\pi^{15}$$
$$-43097853365074859529398834611606304296382250 0\sqrt{3}\pi^{16}+20360909210142211267479945147389977545645312 5\pi^{17}\Bigg],$$

$$\hat{Z}^{(18,20)}(6)=\frac{1}{65245723071090068417518351071738439318237761817804800000\pi^{18}}\Big[259625842126240500100938908160000+111073180950541186268823643055984640000\sqrt{3}\pi$$
$$+233038023732589927256784920292111360000\pi^2-1032472987639798739422860608045285376000\sqrt{3}\pi^3-20666791715085597958599016781040153600000\pi^4$$
$$+350029277432491841213725543224307746816000\sqrt{3}\pi^5+867475371849645978993078753504572219904000\pi^6-671606849547530437668819265336557905510400\sqrt{3}\pi^7$$
$$-193075475826251205514087363454187691555200\pi^8+860772516542746194069386453276313232013568\sqrt{3}\pi^9+2383941001450039629520003136254956831826752\pi^{10}$$
$$-74059628907079159742149540945021561913118720\sqrt{3}\pi^{11}-16261634604298997296661982480767219212620000\pi^{12}+384490645501206647632780617040348429495701888\sqrt{3}\pi^{13}$$
$$+55804905927834385668587867717361480497260579\pi^{14}-997256308062756874082224554265091923222520832\sqrt{3}\pi^{15}-654657249271827076492025979509844811361784750\pi^{16}$$
$$+867271418676735667061337537804433499754022500\sqrt{3}\pi^{17}-36380959856686791483468353731868758441029687 5\pi^{18}\Big],$$

$$\hat{Z}^{(19,21)}(6)=\frac{1}{2060478203387484265844386431983800000432957624779584569344000000\pi^{19}}\Big[-1590489804299012371497908036081193984000\sqrt{3}+414582078560004945198851894219967759360000\pi$$
$$+22280771287352948633529820055966228244480000\sqrt{3}\pi^2+1563890296669448431813625668700729545344000000\pi^3-140445105490805128670677239365856032074752000\sqrt{3}\pi^4$$
$$-11568091795226820855199468075702123190060697600\pi^5+52014967443522320307594515995415346504041472000\sqrt{3}\pi^6+3543009410059187067520114271434610534471831040000\pi^7$$
$$-11788675047557938287981760697364134875237413120 0\sqrt{3}\pi^8-57731125238403879923103604843172542877582089132 80\pi^9+17979869642561112978013340053037998881868710873 60\sqrt{3}\pi^{10}$$
$$+53078810858202617452122520867312702876287973430400\pi^{11}-18732679458183045964589094996361098775876099382400\sqrt{3}\pi^{12}-26639116503938461697749014553483217362205651540092 8\pi^{13}$$
$$+12354980013319886799027062471455129172746385456505 6\sqrt{3}\pi^{14}+62503414080945722788305827436942566298837936549456 0\pi^{15}-44484845081926181697778758656582409206944666287301 2\sqrt{3}\pi^{16}$$
$$-35733815419750421304108062256926068151449145949577 5\pi^{17}+63050631012298260498912219008644137048091431440625 0\sqrt{3}\pi^{18}-29339428865314237189040857066646761998302423710937 5\pi^{19}\Big],$$

$$\hat{Z}^{(20,22)}(6)=\frac{1}{131870605016798993014040731646963200027709287985893412438016000000\pi^{20}}\Big[5466037859721646893685119321724723200 00+41921680565297237837902160585230834237440 00\sqrt{3}\pi$$
$$+94822995728253394754441178272014676043840000\pi^2-49810869135289421108344964835821556242841600 0\sqrt{3}\pi^3-10813353236658017394381262369568790210769920 00\pi^4$$
$$+20276162513157527117918998975210781124786585600\sqrt{3}\pi^5+62497517617213410112927515282728748416219136000\pi^6-46206743452979637609512743779032844933179965440 0\sqrt{3}\pi^7$$
$$-19598975672972797705915965955325003915881929984 00\pi^8+75754013920491549303243191068914767072117002444 80\sqrt{3}\pi^9+35201870642033135407771614064221957353311528780800\pi^{10}$$
$$-96286971563338005690087005676610320839863334584320\sqrt{3}\pi^{11}-37261759422761595255950131484568989128779394236864 0\pi^{12}+85503370312175830438517572832704665100141749626572 8\sqrt{3}\pi^{13}$$
$$+22818132316639652808573230999733233198591582627968 0\pi^{14}-45211001661327134162372202771205418189801885694136 32\sqrt{3}\pi^{15}-72887087969396587709522985736169826954875673050211 12\pi^{16}$$
$$+11761074955706242770234659317182460615849362412096800\sqrt{3}\pi^{17}+82435109034180846412522527045344103070816050780450 00\pi^{18}$$
$$-10204293076060846374166557075101543051605762249920000\sqrt{3}\pi^{19}+42312859889154424092707005632230672566253794698593 75\pi^{20}\Big],$$

$$\hat{Z}^{(1,4)}(6)=\frac{5}{12\pi}-\frac{2}{9\sqrt{3}},\qquad \hat{Z}^{(2,5)}(6)=-\frac{1728+192\sqrt{3}\pi-281\pi^2}{31104\pi^2},\qquad \hat{Z}^{(3,6)}(6)=\frac{-83592+10368\sqrt{3}\pi+127035\pi^2-22840\sqrt{3}\pi^3}{10077696\pi^3},$$

$$\hat{Z}^{(4,7)}(6)=\frac{4455648+414720\sqrt{3}\pi-11109312\pi^2-971904\sqrt{3}\pi^3+1592539\pi^4}{5804752896\pi^4},\qquad \hat{Z}^{(5,8)}(6)=\frac{238295520-21461760\sqrt{3}\pi-1411557840\pi^2+93070080\sqrt{3}\pi^3+1737995139\pi^4-302777720\sqrt{3}\pi^5}{3134566563840\pi^5},$$

$$\hat{Z}^{(6,9)}(6)=-\frac{76892820480+5576325120\sqrt{3}\pi-573013245600\pi^2-1383091200\sqrt{3}\pi^3+1090674751104\pi^4+79621753536\sqrt{3}\pi^5-148557967675\pi^6}{13541327555788800\pi^6},$$

$$\hat{Z}^{(7,10)}(6)=\frac{-31425707738880+2113494405120\sqrt{3}\pi+429913865757600\pi^2+7801923628800\sqrt{3}\pi^3-1742440664795544\pi^4+32206476790656\sqrt{3}\pi^5+1964049821256195\pi^6-332651386503800\sqrt{3}\pi^7}{716607054252343296000\pi^7},$$

$$\hat{Z}^{(8,11)}(6)=\frac{1}{8255313264986994769920 0\pi^8}\Big[2315467859604480+130161909104640\sqrt{3}\pi-34843187276513280\pi^2+3014457537208320\sqrt{3}\pi^3+177112509501098304\pi^4-11881252279830528\sqrt{3}\pi^5$$
$$-3102522041786768 64\pi^6-19421735763949824\sqrt{3}\pi^7+41007215197066475\pi^8\Big],$$

$$\hat{Z}^{(9,12)}(6)=-\frac{1}{8024164493567358916362240 0\pi^9}\Big[-145843276422251520+7480839015014400\sqrt{3}\pi+36716057881629235 20\pi^2+351856049737973760\sqrt{3}\pi^3-31386652836855414336\pi^4$$
$$-3629211613868090880\sqrt{3}\pi^5+10694870152629355296\pi^6+2180932927606245888\sqrt{3}\pi^7-11522467124695902959 1\pi^8+19058564900547151000\sqrt{3}\pi^9\Big],$$

$$\hat{Z}^{(10,13)}(6)=-\frac{1}{5777398435368498419780812800 00\pi^{10}}\Big[594246584051269632 00+2607345528496128000\sqrt{3}\pi-14946468433496812800 00\pi^2+27671209311139430400 0\sqrt{3}\pi^3+14241443740755440670720\pi^4$$
$$-3145180931422213017600\sqrt{3}\pi^5-6332597588157514581600 0\pi^6+7942059992375119411200\sqrt{3}\pi^7+10772532845464012450732 8\pi^8+5879363304117906600000\sqrt{3}\pi^9-13946884924872838124375\pi^{10}\Big],$$

$$\hat{Z}^{(11,14)}(6)=\frac{1}{7549904275339553734969566167040000\pi^{11}}\Big[-4399641826672218255360 0+17503339917763510272 00\sqrt{3}\pi+17949479284159163516160 00\pi^2+31558521644948248473600 0\sqrt{3}\pi^3$$
$$-2651912159363688811113216 0\pi^4-67910712046503090341068 80\sqrt{3}\pi^5+18038496114614990782896480 0\pi^6+3698599472379594736462080 0\sqrt{3}\pi^7-55973581800661523593769882 4\pi^8$$
$$-2995434405717947973231091 2\sqrt{3}\pi^9+58826463176273287629777097 5\pi^{10}-95359518807271769754935000\sqrt{3}\pi^{11}\Big],$$

$$\hat{Z}^{(12,15)}(6)=\frac{1}{1304623458778674885402741033664512000 0\pi^{12}}\Big[38883507875801345433600 00+13442441672779463393280 0\sqrt{3}\pi-14590689613676603996405760 0\pi^2+44000698719609952911360000\sqrt{3}\pi^3$$
$$+2186295711339895922635545600\pi^4-88513345480448125807460352 0\sqrt{3}\pi^5-17675240499320755967078031360\pi^6+60563947267851870259169280 00\sqrt{3}\pi^7+74493064178295232470870149280\pi^8$$
$$-1312221999574993409584785100 8\sqrt{3}\pi^9-12567387966727966910617210502 4\pi^{10}-60468751117060347876862176 00\sqrt{3}\pi^{11}+16019905458889520591171759375\pi^{12}\Big],$$

$$\hat{Z}^{(13,16)}(6)=-\frac{1}{2381198736962837400837082934644467302400 00\pi^{13}}\Big[-36081029421958636931751936 00+11287958287395384669634560 0\sqrt{3}\pi+22005253339637263910189015040 0\pi^2$$
$$+57827617497511962505858252800\sqrt{3}\pi^3-50057777253066382392304032921 60\pi^4-20880064273599764032760639078 40\sqrt{3}\pi^5+56245971637991179160763170135040\pi^6$$
$$+23616031401582363829127165460480\sqrt{3}\pi^7-33758584027422690667484217901622 4\pi^8-97993822616708448714474411277056\sqrt{3}\pi^9+99412355205780663340477841052657 6\pi^{10}$$
$$+82135186115141484988489908686592\sqrt{3}\pi^{11}-10305906463055375826931116946266 75\pi^{12}+16417299673667795736003277607 5000\sqrt{3}\pi^{13}\Big],$$

$$\hat{Z}^{(14,17)}(6)=-\frac{1}{33603476576019561400612914373702722571468800000\pi^{14}}\Big[2389057014651126852082021171200+65698433766906751948161024000\sqrt{3}\pi-12378106468218687742923121704960\pi^{2}$$
$$+55046198799925296179864430182400\sqrt{3}\pi^{3}+260156842900013267347575599235072\pi^{4}-16699682548006255663567388983296\sqrt{3}\pi^{5}-3254045055645008055641246400139776\pi^{6}$$
$$+2008109501687376242875158200745984\sqrt{3}\pi^{7}+247041181687428123019584262350455808\pi^{8}-1120389330458275127264709547563110\sqrt{3}\pi^{9}-10188996848188880461041893780860856\pi^{10}$$
$$+22529710567670830129065904525689753\sqrt{3}\pi^{11}+17208329668233956283849226303993113\pi^{12}+735699017309472624745927140281808\sqrt{3}\pi^{13}-21667427530753248514433495502232562\pi^{14}\Big],$$

$$\hat{Z}^{(15,18)}(6)=\frac{1}{272188160265758447344964606426992052828897280000\pi^{15}}\Big[-89870038309643918511764894515200+22368420103478267720549007360000\sqrt{3}\pi$$
$$+77531309401121148711700384542720000\pi^{2}+27892526887178816642101198356480000\sqrt{3}\pi^{3}-25145855167835911447345817957234688\pi^{4}-15343889012024235503574964512522240\sqrt{3}\pi^{5}$$
$$+41581557778769108504531140550689152000\pi^{6}+28888856344525009970811969977730048\sqrt{3}\pi^{7}-40163699368365453914272684146196919424\pi^{8}$$
$$-23998369622688852629411845339471964160\sqrt{3}\pi^{9}+22567408468131446617187257231678440408\pi^{10}+84401003720904437598197984963889045120\sqrt{3}\pi^{11}$$
$$-64639819881235469795017534462146301521\pi^{12}-70316314479908100041829924589063018560\sqrt{3}\pi^{13}$$
$$+66543344063591273528809654944904072668\pi^{14}-10439011765691323519628082535885258750\sqrt{3}\pi^{15}\Big],$$

$$\hat{Z}^{(16,19)}(6)=\frac{1}{627121521252307462682798453207789689717779333120000\pi^{16}}\Big[90127565300227557177080441339904000+19892191805627448373193610362880\sqrt{3}\pi$$
$$-60868794617475995242988611043328000\pi^{2}+380615176638169748196166720290816000\sqrt{3}\pi^{3}+16375985035042776078266039041532559360\pi^{4}$$
$$-15865402697558606548012186642132500480\sqrt{3}\pi^{5}-28654135372035185045660954323476480000\pi^{6}+28421999916764922473207798974428610560\sqrt{3}\pi^{7}$$
$$+33964951775142977349118882712190993070080\pi^{8}-27502499397548699839371434246254034288640\sqrt{3}\pi^{9}-25124627321994451350488160944016694886400\pi^{10}$$
$$+13878154255956432555613929901342135050240\sqrt{3}\pi^{11}+10268810213211011184055952474897485946830080\pi^{12}-26845239416616753309667288717073828097638\sqrt{3}\pi^{13}$$
$$-17439423696386539666659308618399194843968\pi^{14}-66621085645103713260100911501297756480000\sqrt{3}\pi^{15}+21737049206082005904134047034810050942187\pi^{16}\Big],$$

$$\hat{Z}^{(17,20)}(6)=-\frac{1}{19573716921327020525255505321521531795471328545341440000\pi^{17}}\Big[-12088923095860164272907930203888025600+24178618278495222838980554816225280\sqrt{3}\pi$$
$$+14177644320711654416353827568445816832000\pi^{2}+6646979543701355919426306325767782400000\sqrt{3}\pi^{3}-619756855153875209831179686368838362726400\pi^{4}$$
$$-52524303073386147285481231854839574036480\sqrt{3}\pi^{5}+13844946268419810027656083744161646898380800\pi^{6}+14838581561193547881840059784280432312320000\sqrt{3}\pi^{7}$$
$$-19121558131230911455820566339830336310582272\pi^{8}-20094119947587557464483578736286521912172544\sqrt{3}\pi^{9}+17198523966992947612465031540569222911359078\pi^{10}$$
$$+13756603625809082794214696882245629685063680\sqrt{3}\pi^{11}-94168632880025613624722595705348336662067083\pi^{12}-43131055976160629703171477969814681559349565\sqrt{3}\pi^{13}$$
$$+26662903847476393636552341568283318328033812544\pi^{14}+35380671670078353128860116073109864925285840\sqrt{3}\pi^{15}$$
$$-27373374398914281806076421955274149565839184375\pi^{16}+42359310415653213327888488169668169689713750\sqrt{3}\pi^{17}\Big],$$

$$\hat{Z}^{(18,21)}(6)=-\frac{1}{5073507426007963720146226979338381041386168358952501248000000\pi^{18}}\Big[1276270449216576216756604567867411660800+22794021436803611611130720023058841600\sqrt{3}\pi$$
$$-10730191048748447800792247310048987709440000\pi^2+9201299212009050401159417516187640135680000\sqrt{3}\pi^3+337332175661691499722478273305492027604992004\pi^4$$
$$-49295388650125071812886895378229386385817600\sqrt{3}\pi^5-76806109355048011111597056147288590587330560\pi^6+11884043452529451179377494650125542088507392007\sqrt{3}\pi^7$$
$$+13312804959484884601524704234134056446640783360\pi^8-16953234735035511015931927281485577730643066880\sqrt{3}\pi^9-15475266592086621668053067406757342878211611648\pi^{10}$$
$$+14835603094300680349329857917026754354304188416\sqrt{3}\pi^{11}+1120045834395682981662352660022192105893173866496\pi^{12}-7140131566821879077771830114481988291006141972487\sqrt{3}\pi^{13}$$
$$-45512580496717779653846022247730092325390208449284\pi^{14}+13555302901698417767739202998808334981536829890567\sqrt{3}\pi^{15}+77882190210628346434914559892373380742180021840007\pi^{16}$$
$$+26692300946165017534399856576542126715597124000\sqrt{3}\pi^{17}-96246417622845005661752165896697010279245804687\pi^{18}\Big],$$

$$\hat{Z}^{(19,22)}(6)=\frac{1}{19780590752519848952106109747044480004156393197884011865702400000\pi^{19}}\Big[-19981123334801427324621351891364947689472000+32394599952602204337194311099411857408000\sqrt{3}\pi$$
$$+30927894755415360893688010970176617456599040\pi^2+18294764911826395561342157366069346026127360\sqrt{3}\pi^3-17430843423186016610125965329643862532704174080\pi^4$$
$$-19945480329497964545862927554041499378044108800\sqrt{3}\pi^5+48700144015095697127876799183746002666584047616\pi^6+79352666928841372060964830269164007188167655424\sqrt{3}\pi^7$$
$$-86749663392106079559474492695577178206656175882240\pi^8-15843373864555574464500886465688625192890976829440\sqrt{3}\pi^9+11143160678864464417253135661331125780335899923148\pi^{10}$$
$$+17383343424094630589964598507628301811234356871987\sqrt{3}\pi^{11}-99856843850012312968536712122705445964433765496535\pi^{12}-10387869124326065679191717460222243658668751539978\sqrt{3}\pi^{13}$$
$$+54972723791046308381814328130595618554038458142739328\pi^{14}+29890762330352061454223597628253936614966067204764672\sqrt{3}\pi^{15}-15563045451071990287283853134620540389950045877529792\pi^{16}$$
$$-24098057729919583688588685887945453548598105132822400\sqrt{3}\pi^{17}+15980055002592426245529447858267930256258347839493187\pi^{18}-24426064582146572034331657158127984482336945966875000\sqrt{3}\pi^{19}\Big],$$

$$\hat{Z}^{(20,23)}(6)=\frac{1}{28484050683628582491032798035744051205985206204952977086611456000000000\pi^{20}}\Big[1104239932025099997773836052462306511028224000+16077974330344863581740132999842087567360000\sqrt{3}\pi$$
$$-11112409193399646958071221383908855197466624000\pi^2+1290387344237375919029063587896855678599168000\sqrt{3}\pi^3+3622724767294202557800593640476737991585287372800\pi^4$$
$$-84145049993118302211708791398228708747389173760000\sqrt{3}\pi^5-10168726634239272227979856057741157569011489177600\pi^6+25148416165209203555714549571809368838453801779200\sqrt{3}\pi^7$$
$$+26230586759198915239786619932679505826191812213145600\pi^8-47382596771754379856743909009782451680354056798208000\sqrt{3}\pi^9-45658935557603555587301692770219266881019502816460800\pi^{10}$$
$$+61621356243332052339939139234212859984868899091001600\sqrt{3}\pi^{11}+50339804450325861088464438616821037642454413100344627200\pi^{12}-52349279194195612451909405853716339390333569029931008\sqrt{3}\pi^{13}$$
$$-35129675227123350335845131292774314691392777548145868800\pi^{14}+24850467532183735592099483897155331294239998303605145600\sqrt{3}\pi^{15}+14170478120689945194103991890942466826148269659825488022\pi^{16}$$
$$-46891482293862283956508791767367882552728431680105139200\sqrt{3}\pi^{17}-24453476857979107854382154413473321177081861831590364000\pi^{18}$$
$$-75400693159663673539329521932937005258200393359700000000\sqrt{3}\pi^{19}+29996619925240338939317620477337462628372360432030859375\pi^{20}\Big].$$

References

[1] K. Becker, M. Becker, and A. Strominger, *Five-branes, membranes and nonperturbative string theory*, *Nucl.Phys.* **B456** (1995) 130–152, [hep-th/9507158].

[2] Y. Hatsuda, M. Marino, S. Moriyama, and K. Okuyama, *Non-perturbative effects and the refined topological string*, arXiv:1306.1734.

[3] J. M. Maldacena, *The Large N limit of superconformal field theories and supergravity*, *Adv.Theor.Math.Phys.* **2** (1998) 231–252, [hep-th/9711200].

[4] O. Aharony, O. Bergman, D. L. Jafferis, and J. Maldacena, *N=6 superconformal Chern-Simons-matter theories, M2-branes and their gravity duals*, *JHEP* **0810** (2008) 091, [arXiv:0806.1218].

[5] V. Pestun, *Localization of gauge theory on a four-sphere and supersymmetric Wilson loops*, *Commun.Math.Phys.* **313** (2012) 71–129, [arXiv:0712.2824].

[6] A. Kapustin, B. Willett, and I. Yaakov, *Exact Results for Wilson Loops in Superconformal Chern-Simons Theories with Matter*, *JHEP* **1003** (2010) 089, [arXiv:0909.4559].

[7] M. Marino and P. Putrov, *Exact Results in ABJM Theory from Topological Strings*, *JHEP* **1006** (2010) 011, [arXiv:0912.3074].

[8] N. Drukker, M. Marino, and P. Putrov, *From weak to strong coupling in ABJM theory*, *Commun.Math.Phys.* **306** (2011) 511–563, [arXiv:1007.3837].

[9] C. P. Herzog, I. R. Klebanov, S. S. Pufu, and T. Tesileanu, *Multi-Matrix Models and Tri-Sasaki Einstein Spaces*, *Phys.Rev.* **D83** (2011) 046001, [arXiv:1011.5487].

[10] N. Drukker, M. Marino and P. Putrov, "Nonperturbative aspects of ABJM theory," JHEP **1111**, 141 (2011) [arXiv:1103.4844 [hep-th]].

[11] H. Fuji, S. Hirano, and S. Moriyama, *Summing Up All Genus Free Energy of ABJM Matrix Model*, *JHEP* **1108** (2011) 001, [arXiv:1106.4631].

[12] K. Okuyama, *A Note on the Partition Function of ABJM theory on S^3*, *Prog.Theor.Phys.* **127** (2012) 229–242, [arXiv:1110.3555].

[13] M. Marino and P. Putrov, *ABJM theory as a Fermi gas*, *J.Stat.Mech.* **1203** (2012) P03001, [arXiv:1110.4066].

[14] M. Hanada, M. Honda, Y. Honma, J. Nishimura, S. Shiba, and Y. Yoshida, *Numerical studies of the ABJM theory for arbitrary N at arbitrary coupling constant*, *JHEP* **1205** (2012) 121, [arXiv:1202.5300].

[15] Y. Hatsuda, S. Moriyama, and K. Okuyama, *Exact Results on the ABJM Fermi Gas*, *JHEP* **1210** (2012) 020, [arXiv:1207.4283].

[16] P. Putrov and M. Yamazaki, *Exact ABJM Partition Function from TBA*, *Mod.Phys.Lett.* **A27** (2012) 1250200, [arXiv:1207.5066].

[17] Y. Hatsuda, S. Moriyama, and K. Okuyama, *Instanton Effects in ABJM Theory from Fermi Gas Approach*, *JHEP* **1301** (2013) 158, [arXiv:1211.1251].

[18] F. Calvo and M. Marino, *Membrane instantons from a semiclassical TBA*, *JHEP* **1305** (2013) 006, [arXiv:1212.5118].

[19] Y. Hatsuda, S. Moriyama, and K. Okuyama, *Instanton Bound States in ABJM Theory*, *JHEP* **1305** (2013) 054, [arXiv:1301.5184].

[20] J. Kallen and M. Marino, *Instanton effects and quantum spectral curves*, arXiv:1308.6485.

[21] A. Kapustin, B. Willett, and I. Yaakov, *Nonperturbative Tests of Three-Dimensional Dualities*, *JHEP* **1010** (2010) 013, [arXiv:1003.5694].

[22] A. Klemm, M. Marino, M. Schiereck, and M. Soroush, *ABJM Wilson loops in the Fermi gas approach*, arXiv:1207.0611.

[23] A. Grassi, J. Kallen, and M. Marino, *The topological open string wavefunction*, arXiv:1304.6097.

[24] Y. Hatsuda, M. Honda, S. Moriyama, and K. Okuyama, *ABJM Wilson Loops in Arbitrary Representations*, *JHEP* **1310** (2013) 168, [arXiv:1306.4297].

[25] A. Cagnazzo, D. Sorokin, and L. Wulff, *String instanton in AdS(4) x CP**3*, *JHEP* **1005** (2010) 009, [arXiv:0911.5228].

[26] O. Aharony, O. Bergman, and D. L. Jafferis, *Fractional M2-branes*, *JHEP* **0811** (2008) 043, [arXiv:0807.4924].

[27] K. Hosomichi, K.-M. Lee, S. Lee, S. Lee, and J. Park, *N=5,6 Superconformal Chern-Simons Theories and M2-branes on Orbifolds*, *JHEP* **0809** (2008) 002, [arXiv:0806.4977].

[28] C.-M. Chang, S. Minwalla, T. Sharma, and X. Yin, *ABJ Triality: from Higher Spin Fields to Strings*, *J.Phys.* **A46** (2013) 214009, [arXiv:1207.4485].

[29] S. Giombi, S. Minwalla, S. Prakash, S. P. Trivedi, S. R. Wadia, *et. al.*, *Chern-Simons Theory with Vector Fermion Matter*, *Eur.Phys.J.* **C72** (2012) 2112, [arXiv:1110.4386].

[30] S. Matsumoto and S. Moriyama, *ABJ Fractional Brane from ABJM Wilson Loop*, *JHEP* **1403** (2014) 079, [arXiv:1310.8051].

[31] D. L. Jafferis, *The Exact Superconformal R-Symmetry Extremizes Z*, *JHEP* **1205** (2012) 159, [arXiv:1012.3210].

[32] N. Hama, K. Hosomichi, and S. Lee, *Notes on SUSY Gauge Theories on Three-Sphere*, *JHEP* **1103** (2011) 127, [arXiv:1012.3512].

[33] H. Awata, S. Hirano, and M. Shigemori, *The Partition Function of ABJ Theory*, *Prog. Theor. Exp. Phys.* (2013) 053B04, [arXiv:1212.2966].

[34] M. Honda, *Direct derivation of "mirror" ABJ partition function*, *JHEP* **1312** (2013) 046, [arXiv:1310.3126].

[35] N. A. Nekrasov and S. L. Shatashvili, *Quantization of Integrable Systems and Four Dimensional Gauge Theories*, arXiv:0908.4052.

[36] S. Bhattacharyya, A. Grassi, M. Marino, and A. Sen, *A One-Loop Test of Quantum Supergravity*, *Class.Quant.Grav.* **31** (2014) 015012, [arXiv:1210.6057].

[37] T. Kitao, K. Ohta and N. Ohta, "Three-dimensional gauge dynamics from brane configurations with (p,q) - five-brane," Nucl. Phys. B **539**, 79 (1999) [hep-th/9808111].

[38] O. Bergman, A. Hanany, A. Karch and B. Kol, "Branes and supersymmetry breaking in three-dimensional gauge theories," JHEP **9910**, 036 (1999) [hep-th/9908075].

[39] M. Marino, *Chern-Simons theory, matrix integrals, and perturbative three manifold invariants*, *Commun.Math.Phys.* **253** (2004) 25–49, [hep-th/0207096].

[40] M. Aganagic, A. Klemm, M. Marino, and C. Vafa, *Matrix model as a mirror of Chern-Simons theory*, *JHEP* **02** (2004) 010, [hep-th/0211098].

[41] M. Marino, *Chern-Simons theory and topological strings*, *Rev.Mod.Phys.* **77** (2005) 675–720, [hep-th/0406005].

[42] M. Tierz, *Soft matrix models and Chern-Simons partition functions*, *Mod.Phys.Lett.* **A19** (2004) 1365–1378, [hep-th/0212128].

[43] A. Kapustin, B. Willett, and I. Yaakov, *Tests of Seiberg-like Duality in Three Dimensions*, arXiv:1012.4021.

[44] A. Giveon and D. Kutasov, *Seiberg Duality in Chern-Simons Theory*, *Nucl.Phys.* **B812** (2009) 1–11, [arXiv:0808.0360].

[45] B. Willett and I. Yaakov, *N=2 Dualities and Z Extremization in Three Dimensions*, arXiv:1104.0487.

[46] M. Honda and S. Moriyama, *Instanton Effects in Orbifold ABJM Theory*, arXiv:1404.0676.

[47] C. A. Tracy and H. Widom, *Proofs of two conjectures related to the thermodynamic Bethe ansatz*, *Commun.Math.Phys.* **179** (1996) 667–680, [solv-int/9509003].

[48] I. R. Klebanov and A. A. Tseytlin, *Entropy of near extremal black p-branes*, *Nucl.Phys.* **B475** (1996) 164–178, [hep-th/9604089].

[49] M. Marino, *Lectures on localization and matrix models in supersymmetric Chern-Simons-matter theories*, *J.Phys.* **A44** (2011) 463001, [arXiv:1104.0783].

[50] S. Banerjee, R. K. Gupta, and A. Sen, *Logarithmic Corrections to Extremal Black Hole Entropy from Quantum Entropy Function*, *JHEP* **1103** (2011) 147, [arXiv:1005.3044].

[51] S. Banerjee, R. K. Gupta, I. Mandal, and A. Sen, *Logarithmic Corrections to N=4 and N=8 Black Hole Entropy: A One Loop Test of Quantum Gravity*, *JHEP* **1111** (2011) 143, [arXiv:1106.0080].

[52] A. Sen, *Logarithmic Corrections to N=2 Black Hole Entropy: An Infrared Window into the Microstates*, arXiv:1108.3842.

[53] A. Sen, *Logarithmic Corrections to Rotating Extremal Black Hole Entropy in Four and Five Dimensions*, *Gen.Rel.Grav.* **44** (2012) 1947–1991, [arXiv:1109.3706].

[54] A. Sen, *Logarithmic Corrections to Schwarzschild and Other Non-extremal Black Hole Entropy in Different Dimensions*, *JHEP* **1304** (2013) 156, [arXiv:1205.0971].

[55] M. Marino and P. Putrov, "Interacting fermions and N=2 Chern-Simons-matter theories," JHEP **1311**, 199 (2013) [arXiv:1206.6346 [hep-th]].

[56] S. A. Yost, *Supermatrix models*, *Int.J.Mod.Phys.* **A7** (1992) 6105–6120, [hep-th/9111033].

[57] R. Dijkgraaf and C. Vafa, *N=1 supersymmetry, deconstruction, and bosonic gauge theories*, hep-th/0302011.

[58] R. Dijkgraaf, S. Gukov, V. A. Kazakov, and C. Vafa, *Perturbative analysis of gauged matrix models*, *Phys.Rev.* **D68** (2003) 045007, [hep-th/0210238].

[59] R. Gopakumar and C. Vafa, *On the gauge theory / geometry correspondence*, *Adv.Theor.Math.Phys.* **3** (1999) 1415–1443, [hep-th/9811131].

[60] G. Lockhart and C. Vafa, *Superconformal Partition Functions and Non-perturbative Topological Strings*, arXiv:1210.5909.

[61] R. Gopakumar and C. Vafa, *M theory and topological strings. 2.*, hep-th/9812127.

[62] N. A. Nekrasov, *Seiberg-Witten prepotential from instanton counting*, *Adv.Theor.Math.Phys.* **7** (2004) 831–864, [hep-th/0206161].

[63] A. Iqbal, C. Kozcaz, and C. Vafa, *The Refined topological vertex*, *JHEP* **0910** (2009) 069, [hep-th/0701156].

[64] D. Martelli and J. Sparks, *The large N limit of quiver matrix models and Sasaki-Einstein manifolds*, *Phys.Rev.* **D84** (2011) 046008, [arXiv:1102.5289].

[65] S. Cheon, H. Kim, and N. Kim, *Calculating the partition function of N=2 Gauge theories on S^3 and AdS/CFT correspondence*, *JHEP* **1105** (2011) 134, [arXiv:1102.5565].

[66] D. L. Jafferis, I. R. Klebanov, S. S. Pufu, and B. R. Safdi, *Towards the F-Theorem: N=2 Field Theories on the Three-Sphere*, *JHEP* **1106** (2011) 102, [arXiv:1103.1181].

[67] D. R. Gulotta, C. P. Herzog, and S. S. Pufu, *From Necklace Quivers to the F-theorem, Operator Counting, and T(U(N))*, *JHEP* **1112** (2011) 077, [arXiv:1105.2817].

[68] D. R. Gulotta, J. Ang, and C. P. Herzog, *Matrix Models for Supersymmetric Chern-Simons Theories with an ADE Classification*, *JHEP* **1201** (2012) 132, [arXiv:1111.1744].

[69] D. R. Gulotta, C. P. Herzog, and T. Nishioka, *The ABCDEF's of Matrix Models for Supersymmetric Chern-Simons Theories*, *JHEP* **1204** (2012) 138, [arXiv:1201.6360].

[70] R. C. Santamaria, M. Marino, and P. Putrov, *Unquenched flavor and tropical geometry in strongly coupled Chern-Simons-matter theories*, *JHEP* **1110** (2011) 139, [arXiv:1011.6281].

[71] T. Suyama, *Eigenvalue Distributions in Matrix Models for Chern-Simons-matter Theories*, *Nucl.Phys.* **B856** (2012) 497–527, [arXiv:1106.3147].

[72] T. Suyama, *On Large N Solution of N=3 Chern-Simons-adjoint Theories*, *Nucl.Phys.* **B867** (2013) 887–912, [arXiv:1208.2096].

[73] T. Suyama, *A Systematic Study on Matrix Models for Chern-Simons-matter Theories*, *Nucl. Phys.* **B 874** (2013) 528–566, [arXiv:1304.7831].

[74] M. Mezei and S. S. Pufu, *Three-sphere free energy for classical gauge groups*, *JHEP* **1402** (2014) 037, [arXiv:1312.0920].

[75] A. Grassi and M. Marino, *M-theoretic matrix models*, arXiv:1403.4276.